Kansas Agricultural Experiment Station Research Reports

Volume 4
Issue 3 *Southeast Agricultural Research Center
Reports*

Article 9

2018

Timing of Side-Dress Applications of Nitrogen for Corn in Conventional and No-Till Systems

D. W. Sweeney
Kansas State University, dsweeney@ksu.edu

D. E. Shoup
Kansas State University, dshoup@ksu.edu

Dorivar Ruiz Diaz
Kansas State University, ruizdiaz@ksu.edu

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Part of the [Agronomy and Crop Sciences Commons](#)

Recommended Citation

Sweeney, D. W.; Shoup, D. E.; and Ruiz Diaz, Dorivar (2018) "Timing of Side-Dress Applications of Nitrogen for Corn in Conventional and No-Till Systems," *Kansas Agricultural Experiment Station Research Reports*: Vol. 4: Iss. 3. <https://doi.org/10.4148/2378-5977.7570>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2018 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.

Timing of Side-Dress Applications of Nitrogen for Corn in Conventional and No-Till Systems

D.W. Sweeney, D. Shoup, and D. Ruiz-Diaz¹

Summary

Corn yield and yield components were affected by tillage and nitrogen (N) side-dress options in 2017. Corn yields were 14% greater with conventional tillage than with no-till. Yields were improved by either splitting N rate between pre-plant and side-dress or adding additional side-dress N as compared with applying 150 lb/a pre-plant. Side-dress applications of 50 lb N/a at V10 following 150 lb/a applied pre-plant resulted in greatest corn yield.

Introduction

Environmental conditions vary widely in the spring in southeastern Kansas. As a result, much of the N applied prior to corn planting may be lost before the time of maximum plant N uptake. Side-dress or split applications to provide N during rapid growth periods may improve N use efficiency while reducing potential losses to the environment. The objective of this study was to determine the effect of timing of side-dress N fertilization compared with pre-plant N applications for corn grown on a claypan soil.

Experimental Procedures

The experiment was established in spring 2015 on a Parsons silt loam soil at the Parsons unit of the Kansas State University Southeast Agricultural Research Center. The experiment was a split-plot arrangement of a randomized complete block design with four blocks (replications). Whole plot tillage treatments were conventional tillage (chisel, disk, and field cultivate) and no tillage. Sub-plot nitrogen treatments were six pre-plant/side-dress N application combinations that include 1) a no-N control, 2) 150 lb N/a applied pre-plant, 3) 100 lb N/a applied pre-plant with 50 lb N/a applied at the V6 (six-leaf) growth stage, 4) 100 lb N/a applied pre-plant with 50 lb N/a applied at the V10 (ten-leaf) growth stage, 5) 150 lb N/a applied pre-plant with 50 lb N/a applied at the V6 growth stage, and 6) 150 lb N/a applied pre-plant with 50 lb N/a applied at the V10 growth stage. The N source for all treatments was liquid urea-ammonium nitrate (28% N) fertilizer. Pre-plant N fertilizer was applied on March 16, 2017, side-dress N at V6 on May 25, 2017, and side-dress N at V10 on June 12, 2017, to appropriate plots.

¹Department of Agronomy, Kansas State University, Manhattan, KS.

All N was broadcast applied with 7-stream pattern fertilizer nozzles. Corn was planted on April 11 and harvested on September 11, 2017.

Results and Discussion

In 2017, corn yielded 18 bu/a more with conventional tillage than with no-tillage, likely because of 16% greater stand (Table 1). Adding N fertilizer, generally, more than doubled yields obtained in the no-N control. Splitting the N fertilizer to apply 100 lb N/a preplant followed by 50 lb N/a at the V6 or V10 growth stages improved yields by more than 15 bu/a greater than all N applied pre-plant. Adding 50 lb N/a extra at the V6 growth stage to a 150 lb N/a preplant application did not improve yields more than that obtained with 150 lb N/a applied split pre-plant and side-dress. However, delaying the extra 50 lb N/a side-dress application to the V10 stage improved yield by nearly 20 bu/a. These effects of N timing on corn yield in 2017 appeared to be related to the combined responses in kernel weight, ears/plant and kernels/ear.

Table 1. Tillage and nitrogen (N) side-dress application effects on yield and yield components of corn in 2017

Treatment	Yield bu/a	Stand number/a	Kernel weight mg	Ears/plant	Kernels/ear
Tillage					
Conventional ¹	147.3	22300	225	0.93	789
No-till	129.0	19200	230	0.90	800
LSD (0.10)	16.6	1300	NS	NS	NS
N timing ²					
No-N control	56.1	20900	178	0.82	483
150 PP	134.8	20900	220	0.92	814
100 PP/50 V6	152.0	20500	232	0.95	866
100 PP/50 V10	151.1	20600	240	0.92	850
150 PP/50 V6	157.8	20800	246	0.96	826
150 PP/50 V10	177.0	20900	250	0.94	929
LSD (0.05)	15.2	NS	19	0.08	80

¹Conventional tillage: chisel, disk, and field cultivate.

²Nitrogen treatments: Control, no N fertilizer; 150 PP, 150 lb N/a applied pre-plant with no side-dress N; 100 PP/50 V6, 100 lb N/a applied pre-plant with 50 lb N/a side-dress applied at V6 (six-leaf) growth stage; 100 PP/50 V10, 100 lb N/a applied pre-plant with 50 lb N/a side-dress applied at V10 (ten-leaf) growth stage; 150 PP/50 V6, 150 lb N/a applied pre-plant with 50 lb N/a side-dress applied at V6 growth stage; and 150 PP/50 V10, 150 lb N/a applied pre-plant with 50 lb N/a side-dress applied at V10 growth stage.