Kansas Agricultural Experiment Station Research Reports

Volume 4
Issue 6 *Turfgrass Research*

Article 5

2018

Evaluating Zoysiagrass-Tall Fescue Mixtures in Kansas

Mingying Xiang

Kansas State University, mxiang@ksu.edu

Jack D. Fry

Kansas State University, jfry@ksu.edu

Megan M. Kennelly

Kansas State University, kennelly@ksu.edu

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Part of the [Horticulture Commons](#)

Recommended Citation

Xiang, Mingying; Fry, Jack D.; and Kennelly, Megan M. (2018) "Evaluating Zoysiagrass-Tall Fescue Mixtures in Kansas," *Kansas Agricultural Experiment Station Research Reports*: Vol. 4: Iss. 6. <https://doi.org/10.4148/2378-5977.7595>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2018 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.

Evaluating Zoysiagrass-Tall Fescue Mixtures in Kansas

Cover Page Footnote

Contribution number 18-630-S.

TURFGRASS RESEARCH 2018

JULY 2018

K-STATE
Research and Extension

Kansas State University
Agricultural Experiment Station
and Cooperative Extension Service

K-State Research and Extension is an equal
opportunity provider and employer.

Evaluating Zoysiagrass-Tall Fescue Mixtures in Kansas

Mingying Xiang, Jack D. Fry, and Megan M. Kennelly

Summary

Water conservation is increasingly important when selecting turfgrasses. Zoysiagrass (*Zoysia japonica*), a C4 grass, is more drought resistant than C3 grasses. However, there is some resistance to the use of zoysiagrass in lawns and golf courses due to its extended dormant period when turf is brown. The objective of this study was to evaluate mowing height, and tall fescue seeding rate and time for establishment of a mixed stand with seeded zoysiagrass. Seeding zoysiagrass and tall fescue together in June generally resulted in a less uniform mixture than seeding zoysiagrass in June and then seeding tall fescue in September. Tall fescue seeded in September at 8 lb of pure live seed (PLS)/1,000 ft² resulted the most uniform mixture of the two species.

Rationale

Water conservation is an increasingly important factor when selecting turfgrasses for use in the landscape. Zoysiagrass (*Zoysia japonica*), a warm-season grass, is more drought resistant than cool-season grasses. However, there is resistance on the part of homeowners, and some golf course managers, to use zoysiagrass due to its extended period of brown color when dormant, which is longer than cool-season grasses.

Mixing zoysiagrass with a cool-season grass, such as tall fescue (*Schedonorus arundinaceus*), to establish a perennial sward has several potential advantages. First, tall fescue enhances late fall and early spring color of the sward while zoysiagrass provides quality during periods of drought stress. In addition, a mixture will allow adaptation across a range of environmental conditions and enhanced resistance to biotic pests. Therefore, this mixture may provide a quality sward allowing reduced irrigation, fertilizer, and pesticide inputs compared to monostands of either species. This

View all turfgrass research reports online at: <http://newprairiepress.org/kaesrr>

combination also maintains acceptable visual color in autumn and early spring when zoysiagrass monostands exhibit a brown, dormant color.

Objectives

The objectives of this research were to evaluate the effects of mowing height, and 'Corona' tall fescue seeding time and rate, on establishment of a perennial mixed sward with 'Compadre' zoysiagrass in Kansas.

Study Description

A field study was initiated in July 2015 at the Kansas State University Rocky Ford Turfgrass Research Center in Manhattan and Olathe, KS. For simplicity, only Manhattan data are shown in this report. Plots were arranged in a split-plot design, with mowing height (0.75 or 2 inches, representing fairway and lawn height, respectively) as the whole plot. Sub-plots consisted of timing and rate of tall fescue seeding. Whole plots measured 10 × 15 ft and were replicated four times. Sub plots were 5 × 5 ft and there were six sub-plot replicates within each whole plot. 'Compadre' zoysiagrass was seeded in all plots at 1 lb of PLS/1,000 ft² on June 9, 2015. 'Corona' tall fescue was seeded at differing times and rates as the sub-plot treatments: 1) June, at the same time zoysiagrass was seeded; and 2) in September into the established stand of zoysiagrass. Data were collected on visual turf quality each week on a 1 to 9 scale (1 = poorest color, density, and uniformity; 6 = minimum acceptable quality; 9 = optimum color, density, and uniformity) and fall color (on a 1 to 9 scale, 1 = brown and 9 = dark green). Tall fescue incidence was determined in December, when zoysiagrass had lost all green color, by counting the presence of tall fescue under a 196-intersection grid that measured 4.6 × 4.6 ft with 2.5 inches between each of 14 gridlines in either direction (% tall fescue incidence = tall fescue frequency/196 × 100). Percentage of tall fescue cover in December was also determined using digital image analysis. All data were subjected to analysis of variance using the GLIMMIX procedure of SAS 9.4 (SAS Institute Inc., Cary, NC). Fisher's protected least significant difference (LSD) ($P \leq 0.05$) was used to detect treatment differences.

Results

A significant mowing height × timing × date of seeding interaction and a significant rate × date interaction were observed on tall fescue incidence (%) determined using the grid. Seeding tall fescue in September resulted in higher incidence of this species at both mowing heights (Figure 1). Furthermore, seeding tall fescue at 8 lb of PLS/1,000 ft² provided higher tall fescue incidence compared to other seeding rates (Figure 2), and this resulted in better late fall and early spring color.

K-STATE
Research and Extension

Kansas State University
Agricultural Experiment Station
and Cooperative Extension Service

Figure 1. Effects of mowing height and time of seeding on tall fescue incidence. On each date, means followed by the same letter are not significantly different ($P \leq 0.05$).

K-STATE
Research and Extension

Kansas State University
Agricultural Experiment Station
and Cooperative Extension Service

Figure 2. Effects of seeding rate on tall fescue incidence. Within each date, means with the same letters are not significantly different ($P \leq 0.05$). PLS = pure live seed.

K-STATE
Research and Extension

Kansas State University
Agricultural Experiment Station
and Cooperative Extension Service

Figure 3. Study area at the Rocky Ford Research Center in Manhattan, KS, on December 4, 2015.

Top image: A-F at 0.75 inches mowing height. All plots seeded with 1 lb/1,000 ft² zoysia in June. Tall fescue seeding: A) 8 lb/1,000 ft² in September; B) 8 lb/1,000 ft² in June; C) 4 lb/1,000 ft² in September; D) 2 lb/1,000 ft² in June; E) 4 lb/1,000 ft² in June; and F) 2 lb/1,000 ft² in September.

Bottom image: G-L at 2 inches mowing height. All plots seeded with 1 lb/1,000 ft² zoysia in June. Tall fescue seeding: G) 2 lb/1,000 ft² in September; H) 2 lb/1,000 ft² in June; I) 8 lb/1,000 ft² in September; J) 1 lb/1,000 ft² zoysia in June; 4 lb/1,000 ft² in June; K) 8 lb/1,000 ft² in June; and L) 4 lb/1,000 ft² in September.

Figure 4. Study area at the Rocky Ford Research Center in Manhattan, KS, on March 31, 2016.

All plots seeded with 1 lb/1,000 ft² zoysia in June. Tall fescue seeding: A) 8 lb/1,000 ft² in June at 0.75 inches mowing height; B) 8 lb/1,000 ft² in September at 0.75 inches mowing height; C) 2 lb/1,000 ft² in September at 0.75 inches mowing height; D) 4 lb/1,000 ft² in September at 0.75 inches mowing height; E) 8 lb/1,000 ft² in September at 2 inches mowing height; and F) 8 lb/1,000 ft² in September at 2 inches mowing height; G) 8 lb/1,000 ft² in September at 0.75 inches mowing height.

K-STATE
Research and Extension

Kansas State University
Agricultural Experiment Station
and Cooperative Extension Service

Figure 5. Study area at the Rocky Ford Research Center in Manhattan, KS, on April 19, 2018.

All plots seeded with 1 lb/1,000 ft² zoysia in June. Tall fescue seeding: A) 8 lb/1,000 ft² in June at 0.75 inches mowing height; B) 8 lb/1,000 ft² in September at 0.75 inches mowing height; C) 2 lb/1,000 ft² in September at 0.75 inches mowing height; D) 4 lb/1,000 ft² in September at 0.75 inches mowing height; E) 8 lb/1,000 ft² in September at 2 inches mowing height; and F) 8 lb/1,000 ft² in September at 2 inches mowing height; G) 8 lb/1,000 ft² in September at 0.75 inches mowing height.

K-STATE
Research and Extension

Kansas State University
Agricultural Experiment Station
and Cooperative Extension Service