Kansas Agricultural Experiment Station Research Reports

Volume 4 Issue 7 Kansas Field Research

Article 5

2018

Best Management Systems to Intensify Soybean Production

G. R. Balboa Kansas State University, balboa@ksu.edu

I. A. Ciampitti Kansas State University, ciampitti@ksu.edu

Follow this and additional works at: https://newprairiepress.org/kaesrr

Part of the Agronomy and Crop Sciences Commons

Recommended Citation

Balboa, G. R. and Ciampitti, I. A. (2018) "Best Management Systems to Intensify Soybean Production," *Kansas Agricultural Experiment Station Research Reports*: Vol. 4: Iss. 7. https://doi.org/10.4148/2378-5977.7606

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2018 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.





KANSAS FIELD RESEARCH 2018



Best Management Systems to Intensify Soybean Production

G.R. Balboa and I.A. Ciampitti

Summary

The aim of this study was to evaluate different management systems to close the yield gap in soybean production. A soybean experiment was established in Scandia, KS, evaluating five management systems under both rainfed and irrigated conditions. For the 2017 season, dryland and irrigated average yields were similar (63–65 bu/a) due to herbicide injury on the irrigated phase. In both water scenarios, intensification (high input) increased yields compared with common practice (low input) systems. Under irrigation, a consistent response to a balanced nutrition program was documented.

Introduction

Yield gap is defined by the difference between potential and actual yield. Management practices such as row spacing, seeding rate, fertilization, pest, and disease control affect the size of the yield gap. A management system is a combination of production practices. The aim of this study was to evaluate the combination of production practices to identify the best management systems (BMSs) for closing soybean yield gaps.

Procedures

A soybean experiment was established during the 2017 season at Scandia, KS. This experiment is part of a long-term corn-soybean rotation. A total of five treatments were established in a randomized completely block design with five replications (Table 1).

Prior to the experiment, soil samples were collected in both water environments and analyzed for organic matter %, pH, and phosphorus (P) content (Table 2).

The weather for the 2017 growing season was compared with the 30 years of data for the Scandia location (Figure 1). Precipitation and mean temperature were below the mean for the April - October period compared to the historical 30-year average. Black dots in Figure 1 represent the precipitation and temperature recorded for the past soybean growing seasons for this study.

Results

Grain Yield

The average yield for the dryland condition was 63 bu/a, ranging from 48 to 76 bu/a (Figure 2). The irrigated soybean average yield was 65 bu/a overall, presenting a range

KANSAS FIELD RESEARCH 2018

from 48 to 77 bu/a. Seasonal precipitation was 16.3 in. and the irrigated phase received 5.3 in. of water.

Herbicide injury on irrigated plots was assessed on July 27, negatively affecting the final yield (Figure 4). The minimum yield registered for irrigated was 48 bu/a for common practices (CP) and the maximum 77 bu/a for the ecological intensification (EI) treatment. The balanced nutrition program (CF) under irrigated conditions yielded 11 bu/a more over CP. Under both water conditions (dryland and irrigated), treatment EI and advanced plus (AD) showed the greatest yields, without statistically differing between them (Figure 2). The CF and PI treatments yielded more than CP under irrigation (by 11 and 18 bu/a, respectively); while in dryland condition, CF was superior in yield over CP (+3 bu/a) but without statistically differing. Production intensification with balanced nutrition (EI, AD) allowed obtaining 55% and 58% more yield than CP for dryland and irrigated conditions, respectively (Figure 2). After four years of rotation in high yielding environments (irrigated), CP yields statistically differed from the rest of the treatments showing the impact of the lack of a balanced nutrition program.

Long-Term Rotation

To explore the long-term impact of the different treatments under dryland and irrigated conditions, average yields for 2014, 2015, 2016 and 2017 growing seasons were summarized (Figure 3). Narrowing rows and increasing the plant population increased yields (average 62 bu/a). The yield level in the dryland environment did not show response to fertilization over control treatment. For the irrigated scenario, the average of four growing seasons is showing larger yield differences between treatments compared to the 2017 season alone (Figures 2 and 3). For this scenario, a balanced nutrition program (CF) on top of common practices (CP) increased yields over control treatments (+7 bu/a). Ecological intensification and advanced plus were the highest yielding treatments at 81 and 78 bu/a, respectively.

The 4-year summary provides a synthesis on the impact of different management practices and water scenarios on soybean yields, and can help to better understand the interaction of production practices to identify the BMSs to intensify soybean yield environments. Overall, intensified management systems based on seeding rate increase, narrow row spacing and a balanced nutrition program increased yields compared to common practices.

	1 /	,			
Treatments	СР	CF	PI	EI	AD
Seeding rate	110,000	110,000	175,000	175,000	175,000
Row spacing (inch)	30	30	15	15	15
Fertilization	No P-K-S	P-K-S	No P-K	P-K-S-N	P-K-S-N
Micronutrients	No	No	No	$1 \times (Fe, Zn, B)$	$2 \times (Fe, Zn, B)$
Fungicide/insecticide	No	No	No	$1 \times$	$2 \times$

Table 1. Treatment description, Scandia, KS

CP = Common practices, CF = comprehensive fertilization, PI = production intensification, EI = ecological intensification (CF+PI), AD = advanced plus. P = phosphorus, K = potassium, S = sulfur, N = nitrogen, Fe = iron, Zn = zinc, B = boron.

Table 2. Soil characterization before planting time

Soybean studies	Organic matter %	pН	Phosphorus (ppm)
Irrigated	2.2	6.2	11.0
Dryland	2.3	5.4	7.4



Figure 1. Yearly (1980 – 2016) mean temperature and mean precipitation for the period April – October. Black circles indicate seasons when experimental data were collected. Empty circles indicate years when experimental data were simulated. Dotted vertical and horizontal lines indicate mean temperature (°F) and mean cumulative precipitation (in.) for the period. Percentage of years in each category are listed in parenthesis.

KANSAS FIELD RESEARCH 2018



Figure 2. Soybean grain yield by treatment for dryland and irrigated conditions, Scandia, KS, 2017. Different letter shows statistical differences (P < 0.05). (1) CP = Common practices, (2) CF = comprehensive fertilization, (3) PI = production intensification, (4) EI = ecological intensification (CF+PI), (5) AD = advanced plus.



Figure 3. Soybean grain yield by treatment for dryland and irrigated conditions, Scandia, KS, 2014 – 2017. Different letter shows statistical differences (*P* < 0.05). (1) CP = Common practices, (2) CF = comprehensive fertilization, (3) PI = production intensification, (4) EI = ecological intensification (CF+PI), (5) AD = advanced plus.

Kansas State University Agricultural Experiment Station and Cooperative Extension Service



Figure 4. Crop injury assessed on July 27 with herbicide application on a contiguous soybean field with irrigated plots, Scandia, KS, 2017.