Kansas Agricultural Experiment Station Research Reports

Volume 5
Issue 2 *Southeast Research and Extension*
Center Agricultural Research

Article 10

2019

Pre-Plant Nitrogen Rate and Application Method and Side-Dress Nitrogen Rate Effects on Corn Grown No-Till on a Claypan Soil

D. W. Sweeney
Kansas State University, dsweeney@ksu.edu

D. Ruiz-Diaz
Kansas State University, ruizdiaz@ksu.edu

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Part of the [Agronomy and Crop Sciences Commons](#)

Recommended Citation

Sweeney, D. W. and Ruiz-Diaz, D. (2019) "Pre-Plant Nitrogen Rate and Application Method and Side-Dress Nitrogen Rate Effects on Corn Grown No-Till on a Claypan Soil," *Kansas Agricultural Experiment Station Research Reports*: Vol. 5: Iss. 2. <https://doi.org/10.4148/2378-5977.7736>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2019 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.

Pre-Plant Nitrogen Rate and Application Method and Side-Dress Nitrogen Rate Effects on Corn Grown No-Till on a Claypan Soil

D.W. Sweeney and D. Ruiz-Diaz¹

Summary

Corn yield in 2018 was increased by about 5 bu/a with knife application of pre-plant nitrogen (N) fertilizer compared with broadcast application. Fertilizing with increasing rates of N applied pre-plant, at side-dress, or both had little effect on yield or yield components of corn in 2018.

Introduction

Environmental conditions vary widely in the spring in southeastern Kansas. As a result, much of the N applied prior to corn planting may be lost before the time of maximum plant N uptake. Pre-plant N application method, pre-plant N rate, and side-dress N rate selection to provide N during rapid growth periods may improve N use efficiency while reducing potential losses to the environment. The objective of this study was to determine the effect of timing of pre-plant and side-dress N fertilization options on corn grown on a claypan soil.

Experimental Procedures

The experiment was established in spring 2018 on a Parsons silt loam soil at the Parsons Unit of the Kansas State University Southeast Research and Extension Center that had been in continuous no-till for more than 10 years. The experiment was a factorial arrangement of a randomized complete block design with four blocks (replications). The two factors were pre-plant N fertilizer placement of broadcast and knife (subsurface band at 4 inches deep) and pre-plant/side-dress N rates of 0-0, 0-150, 100-0, 100-50, 100-100, 150-0, 150-50, 150-100, and 200-0 lb/a. Side-dress applications were broadcast at the V10 growth stage using 7-stream pattern fertilizer nozzles. The N source for all treatments was liquid urea-ammonium nitrate (UAN; 28% N) fertilizer. Pre-plant N fertilizer was applied on March 12, 2018, and side-dress N was applied at V10 on June 4, 2018, to appropriate plots. Corn was planted on April 10 and harvested on August 28, 2018.

¹Department of Agronomy, Kansas State University, Manhattan, KS.

Results and Discussion

Even though individual yield components were not significantly affected by pre-plant N application method, general trends resulted in more than 5 bu/a greater corn yields when N was knife applied rather than broadcast prior to planting (Table 1). In general, applying N at any rate and time resulted in approximately 50% greater corn yield in 2018 than the 75.6 bu/a in the no-N control. However, there were few differences in yield among the eight treatments receiving N fertilizer. For example, general increases in total N applied, as well as applying no N until the V10 growth stage (0-150 lb/a pre-plant/side-dress N rate), had little effect on yield in 2018. Stand was not affected by pre-plant/side-dress N rates, but fertilizing with N increased kernel weight, the number of ears/plant, and the number of kernels/ear compared with corn grown in the no-N control.

Table 1. Pre-plant application method and pre-plant/side-dress nitrogen (N) rates effects on yield and yield components of corn planted no-till on a claypan soil in 2018

Treatment	Yield bu/a	Stand plants/a	Kernel weight mg	Ears/plant	Kernels/ear
Pre-plant N method					
Broadcast	101.1	16600	253	1.07	622
Knife ¹	115.6	17200	249	1.10	634
LSD (0.10)	4.4	NS	NS	NS	NS
Pre-plant/side-dress ²					
N rates (lb/a)					
0-0 (No-N control)	75.6	16700	227	1.00	521
0-150	113.2	16000	263	1.05	675
100-0	110.8	16600	249	1.11	625
100-50	116.8	17500	259	1.06	637
100-100	115.3	17300	256	1.13	638
150-0	114.5	16600	251	1.12	643
150-50	114.3	16800	257	1.14	641
150-100	121.2	16600	254	1.13	665
200-0	111.3	17800	243	1.06	609
LSD (0.05)	9.5	NS	15	0.11	70

¹Knife: subsurface band at 4 inch depth.

²Side-dress applications were made at the V10 growth stage.