Kansas Agricultural Experiment Station Research Reports

Volume 6
Issue 4 *Southeast Research and Extension*
Center Agricultural Research

Article 7

2020

Pre-Plant Nitrogen Rate and Application Method and Side-Dress Nitrogen Rate Effects on No-Till Corn Grown on a Claypan Soil

D. W. Sweeney
Kansas State University, dsweeney@ksu.edu

D. Ruiz Diaz
Kansas State University, ruizdiaz@ksu.edu

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Part of the [Agronomy and Crop Sciences Commons](#)

Recommended Citation

Sweeney, D. W. and Ruiz Diaz, D. (2020) "Pre-Plant Nitrogen Rate and Application Method and Side-Dress Nitrogen Rate Effects on No-Till Corn Grown on a Claypan Soil," *Kansas Agricultural Experiment Station Research Reports*: Vol. 6: Iss. 4. <https://doi.org/10.4148/2378-5977.7909>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2020 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.

Pre-Plant Nitrogen Rate and Application Method and Side-Dress Nitrogen Rate Effects on No-Till Corn Grown on a Claypan Soil

Cover Page Footnote

This work is supported by the U.S. Department of Agriculture National Institute of Food and Agriculture, Hatch project KS00-0104-HA.

Pre-Plant Nitrogen Rate and Application Method and Side-Dress Nitrogen Rate Effects on No-Till Corn Grown on a Claypan Soil

D.W. Sweeney and D. Ruiz-Diaz¹

Summary

Average corn yield in 2019 was increased by 14 bu/a with knife application of pre-plant nitrogen (N) fertilizer compared with broadcast application. Applying N more than doubled yield of corn grown without N. In general, applying side-dress N increased yields compared to yields obtained with only pre-plant applications.

Introduction

Environmental conditions vary widely in the spring in southeastern Kansas. As a result, much of the N applied prior to corn planting may be lost before the time of maximum plant N uptake. Pre-plant N application method, pre-plant N rate, and side-dress N rate selections create opportunities to provide N during rapid growth periods and may improve N use efficiency while reducing potential losses to the environment. The objective of this study was to determine the effect of timing of pre-plant and side-dress N fertilization options on corn grown no-till on a claypan soil.

Experimental Procedures

The experiment was established in spring 2018 on a Parsons silt loam soil at the Parsons Unit of the Kansas State University Southeast Research and Extension Center that had been in continuous no-till for more than 10 years. The experiment was a factorial arrangement of a randomized complete block design with four blocks (replications). The two factors were pre-plant N fertilizer placement of broadcast and knife (subsurface band at 4 inches deep) and pre-plant/side-dress N rates of 0-0, 0-150, 100-0, 100-50, 100-100, 150-0, 150-50, 150-100, and 200-0 lb/a. Side-dress applications were broadcast at the V10 growth stage using 7-stream pattern, fertilizer nozzles dropped to less than a foot above the soil surface. The N source for all treatments was liquid urea-ammonium nitrate (UAN; 28% N) fertilizer. Pre-plant N fertilizer was applied on March 19, 2019, and side-dress N was applied at V10 on June 20, 2019, to appropriate plots. Corn was planted on April 11 and harvested on September 4, 2019.

---

¹Department of Agronomy, Kansas State University, Manhattan, KS.

Results and Discussion

Knife application of the N applied pre-plant resulted in 14 bu/a greater yields than when the pre-plant N was broadcast applied (Table 1). This was partially because of approximately 7% greater number of ears per plant with knifing than with broadcasting. The other yield components were not affected by pre-plant application method ($P = 0.05$). Applying N at any rate and time more than doubled corn yield in 2019 compared to the 84 bu/a yield with the no-N control. In general, applying side-dress N increased yields compared to yields obtained with only pre-plant applications; however, the increase from side-dress appeared greater when the pre-plant N was 100 lb N/a than when the pre-plant N was 150 lb N/a. Increasing total N rate to greater than 100 lb N/a resulted in increased yield regardless of individual rates of pre-plant/side-dress N applications, with few differences in combinations where total N was 150 lb/a or greater. Stand was not affected by pre-plant/side-dress N rates, but fertilizing with N increased kernel weight, the number of ears/plant, and the number of kernels/ear compared with corn grown in the no-N control.

Acknowledgment

This work is supported by the U.S. Department of Agriculture National Institute of Food and Agriculture, Hatch project KS00-0104-HA.

Table 1. Pre-plant application method and pre-plant/side-dress nitrogen (N) rates effects on yield and yield components of corn planted no-till on a claypan soil in 2019

Treatment	Yield bu/a	Stand plants/a	Kernel weight mg	Ears/plant	Kernels/ear
Pre-plant N method					
Broadcast	176	21,700	257	1.17	675
Knife ¹	190	21,400	261	1.25	691
LSD (0.10)	6	NS	NS	0.05	NS
Pre-plant/side-dress ²					
N rates (lb/a)					
0-0 (No-N control)	84	21,000	220	0.91	510
0-150	188	21,300	277	1.11	730
100-0	174	21,900	262	1.15	674
100-50	197	22,200	262	1.20	721
100-100	201	21,100	271	1.34	677
150-0	195	21,800	264	1.26	692
150-50	205	21,700	272	1.29	691
150-100	208	21,800	240	1.33	772
200-0	194	21,200	266	1.29	681
LSD (0.05)	13	NS	20	0.10	67

¹Knife: subsurface band at 4 inch depth.

²Side-dress applications were made at the V10 growth stage.