Kansas Agricultural Experiment Station Research Reports

Volume 6
Issue 4 *Southeast Research and Extension*
Center Agricultural Research

Article 10

2020

Effect of Burning and Tillage Options on Yields in a Continuous Wheat-Double-Crop Soybean Rotation

D. W. Sweeney
Kansas State University, dsweeney@ksu.edu

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Part of the [Agronomy and Crop Sciences Commons](#)

Recommended Citation

Sweeney, D. W. (2020) "Effect of Burning and Tillage Options on Yields in a Continuous Wheat-Double-Crop Soybean Rotation," *Kansas Agricultural Experiment Station Research Reports*: Vol. 6: Iss. 4.
<https://doi.org/10.4148/2378-5977.7912>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2020 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.

Effect of Burning and Tillage Options on Yields in a Continuous Wheat-Double-Crop Soybean Rotation

Cover Page Footnote

This work is supported by the U.S. Department of Agriculture National Institute of Food and Agriculture, Hatch project KS00-0104-HA.

Effect of Burning and Tillage Options on Yields in a Continuous Wheat-Double-Crop Soybean Rotation

D.W. Sweeney

Summary

Double-crop soybean yields during the first two years of this study have not been affected by management of previous wheat straw practices by burning or tillage done before planting. However, by the second year of the study, subsequent wheat yields were 41% greater where the wheat residue had been burned the previous year.

Introduction

Double-cropping of soybeans after wheat is practiced by many producers in southeastern Kansas. Several options exist for dealing with wheat straw residue from the previous crop before planting soybeans. However, the method of managing the residue may affect not only the double-crop soybeans but also the following wheat crop. The objective of this study was to determine the effect of burning or no burning with three tillage options (reduced-till, strip-till, and no-till) on double-crop soybean and subsequent wheat yields.

Experimental Procedures

Six wheat residue management systems for double-crop soybean and the subsequent wheat crop were established in spring 2017. The experiment was a split-plot arrangement of a randomized complete block with three replications. The whole plots were burn and no-burn and the subplots were tillage options of reduced-till, strip-till, and no-till prior to planting the double-crop soybeans. In each year after the soybean harvest, the entire area was disked, field cultivated, fertilized, and planted to wheat. Thus, treatment effects on wheat yield was due to the residual from the residue management treatments for the double-crop soybeans.

Results and Discussion

In both 2017 and 2018, burning or not of wheat straw, or tillage prior to planting, did not affect double-crop soybean yields. In 2018, after one year of a continuous wheat-double-crop soybean rotation, subsequent wheat yields were unaffected by the residual of burn or tillage treatments. However, in 2019 wheat yields were 41% greater where the wheat residue had been burned in 2018, even though wheat yields were unaffected by using reduced-, strip-, or no-tillage to plant the previous double-crop soybeans.

Acknowledgment

This work is supported by the U.S. Department of Agriculture National Institute of Food and Agriculture, Hatch project KS00-0104-HA.

Table 1. Effect of residue management on double-crop soybean and subsequent wheat yields

Residue management ¹	Double-crop soybean yields		Wheat yields	
	2017	2018	2018	2019
	----- bu/a -----			
Burn				
Yes	36.4	33.5	55.4	48.5
No	38.2	38.0	55.4	34.3
LSD (0.05)	NS	NS	NS	10.1
Tillage				
Reduced-till	38.3	33.5	55.2	42.4
Strip-till	36.1	36.6	56.9	40.6
No-till	37.4	37.2	54.2	41.2
LSD (0.05)	NS	NS	NS	NS

¹Residue management effects on wheat yields are the residual following those treatments for the double-crop soybeans in the previous year.