

2020

Determining the Effects of Manganese Source and Level in Diets Containing High Levels of Copper on Growth Performance of Growing-Finishing Pigs

H. R. Kerkaert
Kansas State University, kerkaert@k-state.edu

J. C. Woodworth
Kansas State University, jwoodworth@ksu.edu

J. M. DeRouchey
Kansas State University, Manhattan, jderouch@k-state.edu

See next page for additional authors
Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Kerkaert, H. R.; Woodworth, J. C.; DeRouchey, J. M.; Dritz, S. S.; Tokach, M. D.; Goodband, R. D.; and Manzke, N. E. (2020) "Determining the Effects of Manganese Source and Level in Diets Containing High Levels of Copper on Growth Performance of Growing-Finishing Pigs," *Kansas Agricultural Experiment Station Research Reports*: Vol. 6: Iss. 10. <https://doi.org/10.4148/2378-5977.8000>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2020 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Determining the Effects of Manganese Source and Level in Diets Containing High Levels of Copper on Growth Performance of Growing-Finishing Pigs

Abstract

A total of 1,994 pigs (PIC; 337 × 1050; initially 88.2 lb) were used to determine the effect of manganese (Mn) source and level on finishing pig growth performance. This experiment was a follow-up to an Mn source by level study conducted last year. However, unlike last year's study, in the present study all diets contained 150 ppm added Cu from Cu hydroxychloride (IBC; Micronutrients, Indianapolis, IN). Dietary treatments were arranged in a 2 × 3 factorial with main effects of Mn source (Mn hydroxychloride, IntelliBond M, Micronutrients, Indianapolis, IN; or Mn sulfate, MnSO₄, Eurochem, Veracruz, Mexico), and increasing added Mn concentration (8, 16, and 32 ppm). The trace mineral premix was formulated without added Mn. There were 27 pigs per pen and 12 pens per treatment. Diets were corn-soybean meal-distillers dried grains with solubles-based and were fed in 4 phases. Overall, there was no ($P > 0.10$) Mn source × level interaction observed for average daily gain (ADG), average daily feed intake (ADFI), and feed efficiency (F/G). Pigs fed IBM had increased ($P < 0.05$) final body weight (BW), ADG, and ADFI compared to pigs fed MnSO₄. Pigs fed 16 ppm of Mn tended ($P = 0.088$) to have reduced ADFI when compared to pigs fed 8 and 32 ppm of Mn. In conclusion, there appears to be little benefit in growth performance by feeding more than 8 ppm of added Mn. However, pigs fed IBM had improved growth performance compared with those fed MnSO₄. This response is different than our previous study with identical Mn sources but without high levels of added Cu. Further research is needed to understand why we observed an Mn source difference to Mn hydroxychloride when fed in conjunction with pharmacological levels of Cu on pig growth performance.

Keywords

copper, finishing pig, growth, manganese, manganese hydroxychloride

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Cover Page Footnote

The authors would like to thank Micronutrients USA, LLC, Indianapolis, IN, for providing partial financial support for these studies. The authors would like to thank New Horizon Farms, Pipestone, MN, for providing technical assistance for these studies.

Authors

H. R. Kerkaert, J. C. Woodworth, J. M. DeRouchey, S. S. Dritz, M. D. Tokach, R. D. Goodband, and N. E. Manzke

Determining the Effects of Manganese Source and Level in Diets Containing High Levels of Copper on Growth Performance of Growing-Finishing Pigs^{1,2}

Hayden R. Kerkaert, Jason C. Woodworth, Joel M. DeRouchey, Steve S. Dritz,³ Mike D. Tokach, Robert D. Goodband, and Naiana E. Manzke⁴

Summary

A total of 1,994 pigs (PIC; 337 × 1050; initially 88.2 lb) were used to determine the effect of manganese (Mn) source and level on finishing pig growth performance. This experiment was a follow-up to an Mn source by level study conducted last year. However, unlike last year's study, in the present study all diets contained 150 ppm added Cu from Cu hydroxychloride (IBC; Micronutrients, Indianapolis, IN). Dietary treatments were arranged in a 2 × 3 factorial with main effects of Mn source (Mn hydroxychloride, IntelliBond M, Micronutrients, Indianapolis, IN; or Mn sulfate, MnSO₄, Eurochem, Veracruz, Mexico), and increasing added Mn concentration (8, 16, and 32 ppm). The trace mineral premix was formulated without added Mn. There were 27 pigs per pen and 12 pens per treatment. Diets were corn-soybean meal-distillers dried grains with solubles-based and were fed in 4 phases. Overall, there was no ($P > 0.10$) Mn source × level interaction observed for average daily gain (ADG), average daily feed intake (ADFI), and feed efficiency (F/G). Pigs fed IBM had increased ($P < 0.05$) final body weight (BW), ADG, and ADFI compared to pigs fed MnSO₄. Pigs fed 16 ppm of Mn tended ($P = 0.088$) to have reduced ADFI when compared to pigs fed 8 and 32 ppm of Mn. In conclusion, there appears to be little benefit in growth performance by feeding more than 8 ppm of added Mn. However, pigs fed IBM had improved growth performance compared with those fed MnSO₄. This response is different than our previous study with identical Mn sources but without high levels of added Cu.⁵

¹ The authors would like to thank Micronutrients USA, LLC, Indianapolis, IN, for providing partial financial support for these studies.

² The authors would like to thank New Horizon Farms, Pipestone, MN, for providing technical assistance for these studies.

³ Department of Diagnostic Medicine/Pathobiology, College of Veterinary Medicine, Kansas State University.

⁴ Micronutrients, Indianapolis, IN.

⁵ Kerkaert, H.R., J.C. Woodworth, J.M. DeRouchey, S.S. Dritz, M.D. Tokach, and R.D. Goodband. 2019. Determining the effects of manganese source and level on growth performance, carcass characteristics, and economics of growing-finishing pigs. Kansas Agricultural Experiment Station Research. Reports: Vol. 5 Iss. 8. <https://newprairiepress.org/kaesrr/vol5/iss8/23/>.

Further research is needed to understand why we observed an Mn source difference to Mn hydroxychloride when fed in conjunction with pharmacological levels of Cu on pig growth performance.

Introduction

Manganese is an essential trace mineral that is a key component in carbohydrate, lipid and protein metabolism. According to the NRC,⁶ the quantitative requirement for Mn for nursery and finishing diets ranges from 2 to 4 ppm. Assuming bioavailability is not a concern, many swine diets today meet the NRC⁶ estimated requirement for Mn from the major dietary ingredients before a trace mineral premix is added to the diet. However, due to the unknown bioavailability of the innate Mn in ingredients, swine diets typically contain added Mn through a trace mineral premix. In a survey conducted by Flohr et al.,⁷ swine diet Mn levels were found to be supplemented at as low as 3.3 ppm and as high as 40 ppm throughout the entire finishing period. Therefore, there is a wide discrepancy of Mn supplementation in commercial swine diets.

In a previous study, we evaluated the effects of increasing Mn from MnSO₄ or Mn hydroxychloride (IBM; Micronutrients, Indianapolis, IN).⁵ In that study there was no evidence of differences observed between the two Mn sources; however, these diets did not contain high levels of added Cu. Therefore, the objective of this study was to further determine the effects of increasing dietary levels of Mn and the source of Mn on growth performance of growing-finishing pigs fed diets with high levels of added Cu.

Procedures

The Kansas State University Institutional Animal Care and Use Committee approved the protocols used in this experiment.

Animals and Diets

This study was conducted in two barns at a commercial research-finishing site in southwest Minnesota (New Horizon Farms, Pipestone, MN). Each barn was naturally ventilated and double-curtain-sided with a slatted concrete floor and deep manure storage. Each pen (10 × 18 ft) was equipped with a 5-hole stainless steel dry self-feeder (Thorpe Equipment, Thorpe, WI) and a bowl waterer for *ad libitum* access to feed and water.

Two groups of 972 pigs (1,944 total pigs, PIC 337 × 1050; initial BW 80.2 lb) were used in a 100-d growth trial. Pigs were housed in mixed gender pens with 27 pigs per pen and 12 pens per treatment. Daily feed additions to each pen were achieved by using a robotic feeding system (FeedPro; Feedlogic Corp., Wilmar, MN) able to record feed amounts for individual pens. The treatments were structured as a randomized complete block design and arranged in a 2 × 3 factorial with main effects of Mn source (MnSO₄, Eurochem, Veracruz, Mexico; or Mn hydroxychloride, IBM; IntelliBond M, Micronutrients USA, LLC, Indianapolis, IN) and increasing Mn (8, 16, or 32 ppm). All treatment diets were manufactured at the New Horizon Farms Feed Mill in Pipestone,

⁶ NRC. 2012. Nutrient Requirements of Swine. 11th rev. ed. National Academy Press, Washington, DC.

⁷ Flohr, J. R., J.M. DeRouchey, J.C. Woodworth, M.D. Tokach, R.D. Goodband, and S.S. Dritz. 2016. A survey of current feeding regimens for vitamins and trace minerals in the US swine industry. J. Swine Health Prod. 24:290-303.

MN, and were formulated to meet or exceed NRC⁶ requirement estimates for growing-finishing pigs for their respective weight ranges (Table 1). Diets were fed in meal form and in 4 dietary phases within each experiment.

Experimental diets were corn-soybean meal-DDGS-based and were formulated with premix without Mn, Cu, and Zn. Manganese, Cu, and Zn were added to the diet by hand-made premixes, which were added in place of corn in the diet. Each hand-made addition contributed the desired source of Mn, MnSO₄ or IBM, and Mn level, 8, 16, or 32 ppm to the appropriate treatment, along with hand additions to provide 150 ppm of Cu from IntelliBond C (Micronutrients, Indianapolis, IN) and 80 ppm of Zn from IntelliBond Z (Micronutrients, Indianapolis, IN).

Pigs were weighed approximately every 14 days to determine ADG, ADFI, and F/G. Because of the ongoing outbreak of COVID-19, only final weights were obtained, and no carcass data were collected.

Chemical Analysis

Representative diet samples were obtained from all feeders of each treatment and delivered to the Kansas State University Swine Laboratory, Manhattan, KS, and stored at -20°C until analysis. Samples of the diets were combined within dietary treatment, and two composite samples from each treatment were analyzed in duplicate (Cumberland Valley Analytical Services, Hagerstown, MD). Samples were analyzed for Mn, Cu, and Zn content (Method 985.01; AOAC Int., 2000).⁸

Statistical Analysis

Data were analyzed as a randomized complete block design for one-way ANOVA using the lmer function from the lme4 package in R (version 3.5.1 (2018-07-02), R Foundation for Statistical Computing, Vienna, Austria) with pen considered the experimental unit, BW as blocking factor, and treatment as fixed effect. Predetermined orthogonal contrasts were used to evaluate the interactive effects of Mn source and level. Interactions ($P \leq 0.10$) were evaluated linearly or quadratically within the source. All results were considered significant at $P \leq 0.05$ and marginally significant between $P > 0.05$ and $P \leq 0.10$.

Results

The analyzed dietary Mn, Cu, and Zn were consistent with calculated values used in diet formulation and followed the intended Mn titration additions (Table 2).

High levels of Cu were included in all diets and in the grower period, there was an Mn source \times level interaction (linear, $P = 0.029$) observed for F/G, with F/G improving as Mn increased from IBM, but decreasing with increased Mn from MnSO₄. There was no evidence ($P > 0.10$) for Mn source or Mn level effect on ADG, but ADFI was greater ($P = 0.034$) when Mn was provided by IBM. Pigs fed 16 ppm Mn tended ($P = 0.052$) to have decreased ADFI when compared to those fed 8 or 32 ppm, regardless of source.

⁸ AOAC International. 2000. Official methods of analysis of AOAC International. 17th ed. AOAC Int., Gaithersburg, MD.

In the finisher period, there was an Mn source \times level interaction (linear, $P = 0.039$) for ADG, with ADG improving as supplemental Mn was increased for MnSO_4 but decreasing when Mn was increased for pigs fed IBM. There was no evidence ($P > 0.10$) of difference for Mn level to influence ADFI, but ADFI was greater ($P = 0.049$) for pigs fed Mn from IBM compared to pigs fed Mn from MnSO_4 . There was no evidence of difference ($P > 0.10$) for an Mn source or level effect on F/G in the finisher period.

Overall, there was no evidence ($P > 0.10$) for an Mn source \times level interaction for final body weight or any observed growth responses. Pigs fed Mn provided by IBM had greater ($P < 0.05$) ADG and ADFI and heavier ($P < 0.05$) final BW than pigs fed Mn from MnSO_4 . Regardless of the source, pigs fed 16 ppm of Mn tended (quadratic, $P = 0.088$) to have lower overall ADFI than pigs fed 8 or 32 ppm of Mn. There was no evidence of difference ($P > 0.10$) for Mn source or level effect on F/G.

Discussion

According to the NRC,⁶ the Mn requirement estimate for growing-finishing pigs is 2 to 4 ppm of the diet. However, most of the research for determining the Mn requirement was conducted more than 50 years ago. Due to the unknown bioavailability of Mn from ingredients commonly used in diets, Mn is usually added to swine diets through a trace mineral premix, frequently as MnSO_4 . Manganese hydroxychloride is another source of Mn that can be added to trace mineral premixes for swine diets. Hydroxychloride-based minerals are manufactured through the reaction of hydrochloric acid, high purity forms of metal, and water. The products of this reaction are hydroxychloride crystals that contain the desired metal covalently bonded to chloride and hydroxyl groups. The covalent bonds processed by hydroxychloride minerals reduce the ability for the minerals to react with other components of the diet and potentially improve bioavailability.⁹

In a previous study, there was no main effect of Mn source on any influence growth performance or carcass characteristics; however, in the present study pigs fed supplemental Mn from IBM had increased ADG and ADFI and heavier final BW when compared to pigs fed MnSO_4 . The reason for these inconsistent effects of IBM on growth performance was not clear, but could have been a result of the higher levels of dietary Cu in the present experiment. Pharmacological levels of Cu have been shown to improve growth performance in growing-finishing pigs.¹⁰ Copper is excreted from the body through bile. However, when dietary Mn was increased to 200 ppm in rats, Mercandante et al.¹¹ observed a reduction in Cu levels in the bile, signifying a hepatobiliary metabolism relationship between Mn and Cu. With the potential of improved bioavailability of the Mn hydroxychloride, less Cu may have been excreted, allowing for potentially greater utilization of Cu in pigs being fed Mn hydroxychloride. However, this theory warrants further investigation.

⁹ Cao, J., P. R. Henry, R. Guo, R. A. Holwerda, J. P. Toth, R. C. Littell, R. D. Miles, and C. B. Ammerman. 2000. Chemical characteristics and relative bioavailability of supplemental organic zinc sources for poultry and ruminants. *J. Anim. Sci.* 78:2039-2054. doi:10.2527/2000.7882039x.

¹⁰ Coble, K. F., J. M. DeRouche, M. D. Tokach, S. S. Dritz, R. D. Goodband, J. C. Woodworth, and J. L. Usry. 2017. The effects of copper source and concentration on growth performance, carcass characteristics, and pen cleanliness in finishing pigs. *J. Anim. Sci.* doi:10.2527/jas.2017.1624.

¹¹ Mercandante, C. J., C. Herrera, M. A. Pettiglio, M. L. Foster, L. C. Johnson, D. C. Dorman, and T. B. Bartnikas. 2016. The effect of high dose oral manganese exposure on copper, iron, and zinc levels in rats. *Biometals.* 29:417-422. doi:10.1007/s10534-016-9924-6.

Swine Day 2020

In summary, our results suggest that supplementing growing-finishing diets with greater than 8 ppm of Mn did not lead to any improvements in growth performance. Pigs fed Mn provided by IBM had greater ADG, ADFI, and heavier final BW than pigs fed Mn from MnSO_4 . More research is needed to further understand the potential benefits of Mn hydroxychloride fed in conjunction with high levels of Cu on pig growth performance.

Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. Persons using such products assume responsibility for their use in accordance with current label directions of the manufacturer.

Table 1. Composition of basal diet (as-fed basis)¹

Items	Phase 1	Phase 2	Phase 3	Phase 4
Ingredients, %				
Corn	58.80	66.88	72.51	80.66
Soybean meal (46.5% CP)	26.60	18.77	13.29	15.35
DDGS ²	10.00	10.00	10.00	---
Beef tallow	1.50	1.50	1.50	1.50
Limestone, ground	1.08	1.00	0.95	0.73
Monocalcium phosphate (21% P)	0.90	0.75	0.65	0.75
Salt	0.35	0.35	0.35	0.35
L-Lysine-HCl	0.37	0.39	0.39	0.30
DL-Methionine	0.06	0.03	0.01	0.02
L-Threonine	0.09	0.09	0.10	0.10
L-Tryptophan	0.02	0.03	0.03	0.03
Phytase ³	0.04	0.04	0.04	0.04
Vitamin-trace mineral premix ⁴	0.15	0.15	0.15	0.15
Mn source ⁵	+/-	+/-	+/-	+/-
Calculated analysis				
Standardized ileal digestible amino acids, %				
Lysine	1.15	0.97	0.84	0.79
Isoleucine:lysine	63	61	59	60
Leucine:lysine	140	147	155	147
Methionine:lysine	31	30	29	29
Methionine and cysteine:lysine	55	55	56	56
Threonine:lysine	62	62	64	65
Tryptophan:lysine	19	19	19	20
Valine:lysine	70	70	70	70
Lysine:net energy, g/Mcal	4.62	3.82	3.26	3.05
Net energy, kcal/kg	2,486	2,539	2,574	2,594
Crude protein, %	20.8	17.8	15.6	14.4
Calcium, %	0.73	0.63	0.57	0.52
Standardized total tract digestible phosphorus, %	0.52	0.47	0.41	0.39

¹In Exp. 1, phases 1, 2, 3, and 4 were fed from 76.1 to 125.0 lb, 125.0 to 160.1 lb, 160.1 to 220.0 lb, and 220.0 lb to market, respectively. In Exp. 2, phases 1, 2, 3, and 4 were fed from 88.2 to 108.0 lb, 108.0 to 169.8 lb, 169.8 to 229.9 lb, and 229.9 lb to market, respectively.

²DDGS = dried distillers grains with solubles.

³Optiphos 2000 (Huvepharma Inc. Peachtree City, GA) provided 858.7 units of phytase FTU/kg of diet with an assumed release of 0.12 available P.

⁴Tribasic copper chloride (IntelliBond C, Micronutrients, Indianapolis, IN) provided 150 ppm of Cu.

⁵Mn hydroxychloride (IntelliBond M, Micronutrients, Indianapolis, IN); or Mn sulfate (MnSO₄, Eurochem, Veracruz, Mexico).

Table 2. Chemical analysis of diets (as-fed basis)¹

Mineral, ppm	MnSO ₄ , ppm			IBM, ppm		
	8	16	32	8	16	32
Cu	217	207	194	197	199	206
Mn	34	35	50	33	42	52
Zn	126	132	130	126	131	125

¹Values represent means from 16 composite samples (4 per phase). For each treatment, samples were collected from multiple feeders, blended, subsampled, ground, and analyzed (Cumberland Valley Analytical Services, Hagerstown, MD). IntelliBond M (IBM, Micronutrients, Indianapolis, IN).

Table 3. Interactive effects of Mn source and level on grow-finish pig growth performance¹

Item ^{2,3}	MnSO ₄ , ppm			IBM, ppm			SEM	Source × level, <i>P</i> =	
	8	16	32	8	16	32		Linear	Quadratic
BW, lb									
Initial	88.2	88.2	88.2	88.2	88.2	88.0	0.77	0.821	0.650
Grower	167.3	165.6	164.9	168.2	166.2	167.1	1.20	0.468	0.699
Final	286.4	286.2	286.6	292.3	287.5	289.2	1.32	0.391	0.207
Grower									
ADG, lb	1.96	1.94	1.92	1.98	1.94	1.96	0.013	0.330	0.615
ADFI, lb	4.25	4.17	4.23	4.37	4.25	4.30	0.034	0.578	0.879
F/G	2.160	2.155	2.212	2.203	2.183	2.179	0.0051	0.029	0.652
Finisher									
ADG, lb	2.09	2.12	2.14	2.18	2.14	2.14	0.011	0.039	0.258
ADFI, lb	6.19	6.15	6.24	6.35	6.24	6.28	0.030	0.272	0.782
F/G	2.941	2.907	2.915	2.890	2.915	2.941	0.0041	0.267	0.402
Overall									
ADG, lb	2.05	2.05	2.05	2.09	2.05	2.07	0.007	0.284	0.213
ADFI, lb	5.36	5.31	5.40	5.51	5.40	5.42	0.029	0.236	0.856
F/G	2.625	2.604	2.639	2.618	2.632	2.632	0.0038	0.801	0.332

¹A total of 1,944 pigs (initial BW of 88.2 lb) were used in two groups with 27 pigs per pen and 12 replicates per treatment. Mn sources were Mn sulfate (MnSO₄, Erachem, Veracruz, Mexico) or IntelliBond M (IBM, Micronutrients, Indianapolis, IN).

²BW = body weight. ADG = average daily gain. ADFI = average daily feed intake. F/G = feed efficiency. HCW = hot carcass weight.

³The grower period was from d 0 to d 42 in group 1 and from d 0 to 38 in group 2. The finisher period was from d 42 to 97 in group 1 and from d 38 to 100 in group 2.

Swine Day 2020

Table 4. Main effects of Mn source and level on finishing pig growth performance¹

Item ²	Source		SEM	Probability, <i>P</i> =	Level, mg/lb			SEM	Probability, <i>P</i> =	
	MnSO ₄	IBM			8	16	32		Linear	Quadratic
BW, lb										
Initial	88.2	88.2	0.76	0.683	88.2	88.2	88.2	0.76	0.977	0.920
Grower	166.0	167.1	1.12	0.202	167.8	165.8	166.0	1.14	0.153	0.182
Final	286.4	289.7	1.18	0.011	289.2	286.8	287.7	1.21	0.391	0.201
Grower										
ADG, lb	1.94	1.96	0.0095	0.169	1.97	1.94	1.94	0.0104	0.273	0.221
ADFI, lb	4.22	4.30	0.0287	0.034	4.31	4.21	4.26	0.0301	0.478	0.052
F/G	2.174	2.188	0.0041	0.407	2.179	2.169	2.193	0.0043	0.340	0.319
Finisher										
ADG, lb	2.12	2.15	0.0069	0.049	2.15	2.13	2.13	0.0079	0.650	0.352
ADFI, lb	6.20	6.29	0.0213	0.049	6.27	6.19	6.26	0.0240	0.930	0.140
F/G	2.924	2.915	0.0032	0.846	2.915	2.907	2.933	0.0035	0.612	0.671
Overall										
ADG, lb	2.04	2.07	0.0042	0.009	2.07	2.05	2.06	0.0048	0.309	0.103
ADFI, lb	5.36	5.42	0.0238	0.023	5.44	5.36	5.42	0.0254	0.912	0.088
F/G	2.625	2.625	0.0032	0.865	2.618	2.618	2.639	0.0034	0.809	0.332

¹A total of 1,944 pigs (initial BW of 88.2 lb) were used in two groups with 27 pigs per pen and 12 replicates per treatment. Mn sources were Mn sulfate (MnSO₄, Eurochem, Veracruz, Mexico) or IntelliBond M (IBM, Micronutrients, Indianapolis, IN).

²BW = body weight. ADG = average daily gain. ADFI = average daily feed intake. F/G = feed efficiency. HCW = hot carcass weight.

³The grower period was from d 0 to d 42 in group 1 and from d 0 to 38 in group 2. The finisher period was from d 42 to 97 in group 1 and from d 38 to 100 in group 2.