

2021

Differences in Rumination and Animal Activity Can Be Quantified by Utilizing New Technologies

M. A. Scilacci

Kansas State University, morgan08@k-state.edu

A. J. Tarpoff

Kansas State University, tarpoff@k-state.edu

E. C. Titgemeyer

Kansas State University, Manhattan, etitgeme@k-state.edu

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Beef Science Commons](#)

Recommended Citation

Scilacci, M. A.; Tarpoff, A. J.; Titgemeyer, E. C.; O'Quinn, T. G.; Jennings, T.; Vahl, C. I.; Thomson, D. U.; Hollenbeck, W. R.; and Blasi, D. A. (2021) "Differences in Rumination and Animal Activity Can Be Quantified by Utilizing New Technologies," *Kansas Agricultural Experiment Station Research Reports*: Vol. 7: Iss. 1. <https://doi.org/10.4148/2378-5977.8030>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2021 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Differences in Rumination and Animal Activity Can Be Quantified by Utilizing New Technologies

Abstract

Objective: Utilizing new technologies such as Allflex eSense Ear Tags to (Allflex Livestock Intelligence, Madison, WI) evaluate the impacts of limit feeding a high-energy diet in the backgrounding phase on daily rumination and activity in growing cattle.

Study Description: This project was conducted in conjunction with the performance backgrounding study at the Kansas State University Beef Stocker Unit in 2019. Crossbred heifers (n = 418) were outfitted with an ear tag that measured rumination and activity throughout the day and the 84-day backgrounding study. This data was continuously collected via an onsite antenna, downloaded to a computer, and analyzed in a software program.

The Bottom Line: High-energy, limit-fed cattle ruminate less than high roughage, ad libitum cattle by 45 minutes in this study. There were no detected differences in animal activity.

Keywords

limit feeding, rumination, activity

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Cover Page Footnote

National Cattlemen's Beef Association, Kansas Corn Commission

Authors

M. A. Scilacci, A. J. Tarpoff, E. C. Titgemeyer, T. G. O'Quinn, T. Jennings, C. I. Vahl, D. U. Thomson, W. R. Hollenbeck, and D. A. Blasi

Differences in Rumination and Animal Activity Can Be Quantified by Utilizing New Technologies

M.A. Scilacci, A.J. Tarpoff, E.C. Titgemeyer, T.G. O'Quinn, T. Jennings,¹ C.I. Vahl, D.U. Thomson,² W.R. Hollenbeck, and D.A. Blasi

Abstract

The objective of this research was to evaluate the Allflex eSense Flex ear tag (Allflex Livestock Intelligence, Madison, WI), which measures rumination and activity between two different diets fed to growing calves. One diet was high-energy, limit-fed, while the other was high roughage, fed ad libitum. For this study, 418 crossbred heifers of Texas and New Mexico origin were utilized in an 84-day backgrounding study at the Kansas State University Beef Stocker Unit. Coupled with the software data system called Heat-Time Pro+, it was determined that high-energy, limit-fed cattle ruminated less than the high roughage, ad libitum cattle ($P < 0.01$). Using GoPro cameras, footage was also collected to determine activity and enthusiasm differences between diets during feeding time, but the data were inconclusive.

Introduction

Industry-advancing technologies have given cattle producers new opportunities and ways to monitor the performance and health of their animals. The dairy industry, for example, has seen rapid innovation in cow-health monitoring and heat detection for reproduction in recent years. Beef cattle researchers are developing and evaluating similar applications for health or production in the cow-calf sector, stocker-backgrounder phase, and feedlot stage. The Allflex eSense ear tag (Allflex Livestock Intelligence, Madison, WI) provides perspective into the animal that otherwise would largely be unnoticed. Moreover, it measures rumination and general activity through motions of the ear and the animal's body that have been correlated back to a specific behavior (rumination or activity). As it relates to backgrounders and feedlot operators, health monitoring to proactively avoid nutritional or other disruptions to growth is critical. The aim of this study was to observe ruminal and activity differences in high-energy, limit-fed cattle compared to high roughage, ad libitum cattle.

¹ Allflex Livestock Intelligence, Madison, WI.

² Department of Animal Science, College of Agriculture and Life Sciences, Iowa State University, Ames, IA.

Experimental Procedures

A total of 418 weaned, crossbred heifers (body weight = 615 ± 53 lb) were purchased at auction markets in Texas and New Mexico, assembled at two different farms approximately 90 miles southwest of Amarillo, TX, then shipped 570 miles to the Kansas State University Beef Stocker Unit, Manhattan, KS, on May 28, 2019. The heifers were used in a completely randomized block design, 84-day receiving and growing study. The study evaluated the impact of high-energy, limit-fed diets containing dry-rolled corn and Sweet Bran (Cargill Animal Nutrition, Blair, NE) compared to high roughage diets fed ad libitum on both animal rumination and activity, as well as enthusiasm to approach the bunk upon arrival of the feed wagon. Cattle were randomized by arrival weight and assigned to pens, each containing 13 or 14 heifers. Additionally, each pen was randomly assigned to one of two treatments in a "treatment pair." Each treatment pair consisted of one high-energy, limit-fed pen, and one high roughage, ad libitum pen. There was a total of 32 pens. The high roughage, ad libitum diet and the high-energy, limit-fed diet each were formulated to provide 45 or 60 Mcal of net energy for gain/100 lb of dry matter, respectively. Each heifer was tagged with an Allflex eSense Flex Ear Tag to monitor and measure rumination and general activity. In addition, during the first ten days of the study, GoPro cameras were mounted on the feed wagon and tractor to capture footage of the cattle during the feeding process.

Results and Discussion

The high-energy, limit-fed heifers ruminated 45 minutes less per day, compared to the high roughage, ad libitum heifers ($P < 0.01$), as shown in Figure 1. This may be due to less dry matter consumption and lower roughage content of the high-energy, limit-fed diet. However, activity was not affected by diet ($P = 0.70$), which is shown in Figure 2. Based on observation of the GoPro videos collected, it could not be determined if the high-energy, limit-fed cattle were more enthusiastic at feeding time compared to the high roughage, ad libitum cattle.

Implications

Limit feeding practices decreased total time spent ruminating per day by 45 minutes compared to cattle fed a high roughage diet ad libitum. However, it was not conclusive from our data whether one group was generally more active than the other.

Acknowledgments

National Cattlemen's Beef Association
Kansas Corn Commission

Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. Persons using such products assume responsibility for their use in accordance with current label directions of the manufacturer.

CATTLEMEN'S DAY 2021




Figure 1. Total daily rumination, measured in minutes per day, for both high-energy, limit-fed and high roughage, ad libitum heifers.




Figure 2. Total daily activity, measured in minutes per day, for both high-energy, limit-fed and high roughage, ad libitum heifers.