Kansas Agricultural Experiment Station Research Reports

Volume 7
Issue 10 *Animal Feed and Pet Food Research Report*

Article 20

2021

The Effects of Medium Chain Fatty Acids in Mash and Crumbled Pellet Diets on Growth Performance of Broilers

Gage E. Nichols
Kansas State University, gage500@k-state.edu

Haley K. Wecker
Kansas State University, haley27@k-state.edu

Caitlin E. Evans
Kansas State University, caitlinevans@k-state.edu

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

Part of the [Other Animal Sciences Commons](#), and the [Poultry or Avian Science Commons](#)

Recommended Citation

Nichols, Gage E.; Wecker, Haley K.; Evans, Caitlin E.; Jones, Cassandra K.; Stark, Charles R.; and Paulk, Chad B. (2021) "The Effects of Medium Chain Fatty Acids in Mash and Crumbled Pellet Diets on Growth Performance of Broilers," *Kansas Agricultural Experiment Station Research Reports*: Vol. 7: Iss. 10. <https://doi.org/10.4148/2378-5977.8158>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2021 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.

The Effects of Medium Chain Fatty Acids in Mash and Crumbled Pellet Diets on Growth Performance of Broilers

Authors

Gage E. Nichols, Haley K. Wecker, Caitlin E. Evans, Cassandra K. Jones, Charles R. Stark, and Chad B. Paulk

The Effects of Medium Chain Fatty Acids in Mash and Crumbled Pellet Diets on Growth Performance of Broilers

*Gage E. Nichols, Haley K. Wecker, Caitlin E. Evans,
Cassandra K. Jones,¹ Charles R. Stark, and Chad B. Paulk*

Summary

The objective of this experiment was to determine the effects of medium chain fatty acids (MCFA) in mash and crumbled pellet broiler diets. A total of 400 male chicks (Cobb 500; initial BW 0.092 lb) were housed in 4 Petersime batteries and used in an 18-d study. Treatments were randomly assigned to 80 cages within location block resulting in 8 cages per treatment with 5 chicks per cage at placement. Treatments were arranged in a 2 × 5 factorial with main effects of feed type (mash and crumbled pellet) and 0.5% MCFA inclusion (no inclusion, control; hexanoic acid, C6; octanoic acid, C8; decanoic acid, C10; and dodecanoic acid, C12). Fat inclusions in the diets were equalized using 0.5% soybean oil in the control diet. Prior to crumbling, diets were conditioned at 185°F for approximately 20 s and pelleted (CPM, model CL-5, Crawfordsville, IN) with a 5/32 × 7/8 in. ring die. Dietary treatments were fed for the full duration of the study. There was no evidence of feed form × MCFA interactions. From d 0 to 18, chicks fed pelleted diets had improved ($P < 0.001$) body weight gain (BWG), feed intake (FI), feed conversion ratio (FCR), and final BW compared to those fed mash diets. For the overall treatment period, there was no evidence of a MCFA effect ($P > 0.10$) on broiler performance. Pelleting and crumbling feed improved growth performance regardless of MCFA inclusion. The MCFA inclusion did not positively influence growth performance of broilers.

Introduction

Historically, the poultry industry has utilized various feed grade antibiotics to positively influence broiler health resulting in improved growth performance. Consumer pressure, however, has driven the poultry industry to explore ways to reduce or completely remove antibiotics from their production system. Additionally, the implementation of governmental regulations such as the Veterinary Feed Directive has further necessitated effective antibiotic alternatives. The Veterinary Feed Directive final rule, an amendment to the Animal Drug Availability Act of 1996, provides legal control and oversight into the use of medically important antibiotics in feed. Under this regulation, antibiotics such as tetracycline and penicillin are only to be included in feed under the approval and supervision of a licensed veterinarian. One possible antibiotic alternative is the inclusion of medium chain fatty acids in broiler diets. These include caproic acid (C6:0), caprylic acid (C8:0), capric acid (C10:0), and lauric acid (C12:0). Previous

¹ Department of Animal Science and Industry, College of Agriculture, Kansas State University.

research determined that MCFA have a bactericidal effect by changing the cells permeability, resulting in the loss of the cell membrane's integrity, effectively killing the cell.² However, limited data exist determining the effects of MCFA on growth performance of broilers. Therefore, the objective of this experiment was to evaluate MCFA as a dietary additive in mash and crumbled pellet starter diets for broilers.

Materials and Methods

The Institutional Animal Care and Use Committee at Kansas State University (Manhattan, KS) reviewed and approved the protocols used in this experiment. A total of 400 one-day-old male broilers (initially 0.092 lb; Cobb 500, Cobb-Vantress, Siloam Springs, AR) were used in an 18-d study to determine effects of MCFA inclusion in mash or crumbled pellet diets on broiler growth performance. Broilers were placed in 1 of 4 Petersime batteries with 5 broilers per cage (dimensions, 38.0 × 13.0 in.), balanced by BW. Cages were randomly assigned to 1 of 10 dietary treatments within location block with 8 replicates per treatment. Illumination was provided by fluorescent bulbs for the duration of the experiment. Feed was provided *ad libitum* in a 1-pan feeder (capacity was 4.4 lb) per pen. Water was provided *ad libitum* through water troughs. Body weight and feed intake were recorded on day 12 and 18 post-hatching. The BW gain and feed conversion ratio were calculated for each cage for the respective period. Cage mortality was recorded daily and FCR was corrected accordingly.

Dietary treatments

Medium chain fatty acids (Sigma Aldrich, St. Louis, MO) used in this experiment included hexanoic (C6), octanoic (C8), decanoic (C10), and dodecanoic acid (C12) and were guaranteed ≥ 98% purity. Dietary treatments consisted of a 2 × 5 factorial arrangement of treatments with main effects of diet form (mash and crumbled pellet) and MCFA inclusion (control, 0.5% C6, 0.5% C8, 0.5% C10, and 0.5% C12). All diets were formulated with ingredient nutrient specifications and their respective standardized ileal digestible (SID) amino acid coefficients based on Cobb 500 recommended requirements. Diets were manufactured at the Kansas State University O.H. Kruse Feed Technology Innovation Center. Corn was ground to approximately 700 µm. One master batch of feed (2,000 lb) was made and divided into 5 separate batches of feed (400 lb each). Either 0.5% soybean oil for the control or 0.5% MCFA was mixed into each batch of feed to create dietary treatments. Each batch was then divided, with one half remaining in mash form and the other further processed into a crumbled pellet. Prior to crumbling, diets were conditioned at 185°F for approximately 20 s and pelleted (CPM, model CL-5, Crawfordsville, IN) with a 5/32 × 7/8 in. ring die.

Statistical analysis

Data were analyzed using the GLIMMIX procedure in SAS v. 9.4 (SAS Institute Inc., Cary, NC), with pen as the experimental unit, pen location as the blocking factor and adjusted using Tukey-Kramer multiple comparisons. Results were considered significant if $P \leq 0.05$.

² Kim, S. A. and M. S. Rhee. 2013. Marked Synergistic Bactericidal Effects and Mode of Action of Medium-Chain Fatty Acids in Combination with Organic Acids against *Escherichia coli* 0157:H7. Appl. and Envir. Micro. 79:6552-6560. <https://doi.org/10.1128/AEM.02164-13>.

Results and Discussion

From d 0 to 18, there was no evidence of feed form $\times$ MCFA interactions ($P > 0.328$). From d 0 to 18, broilers fed pelleted diets had improved ($P < 0.001$) BW gain, feed intake, FCR, and final BW compared to those fed mash diets. For the overall treatment period there was no evidence of an MCFA effect ($P > 0.119$) on broiler performance. Under the constraints of the current study, pelleting and crumbling feed improved growth performance regardless of MCFA inclusion. Additionally, MCFA inclusion did not positively influence growth performance. More work is needed to determine the optimal inclusion level and type of MCFA required for improvement of broiler growth performance when exposed to a health challenge.

Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. Persons using such products assume responsibility for their use in accordance with current label directions of the manufacturer.

Table 1. Diet composition (as-fed basis)¹

Item	Control	Treatments
Ingredients, %		
Corn	60.76	60.76
Soybean meal (46.5% CP)	32.50	32.50
Soy oil	2.00	1.50
Calcium carbonate	1.40	1.40
Monocalcium P (21% P)	2.10	2.10
Salt	0.23	0.23
L-Lys HCl	0.15	0.15
DL-Met	0.23	0.23
L-Thr	0.08	0.08
Mineral-vitamin premix ²	0.25	0.25
Sodium bicarbonate	0.20	0.20
Choline chloride	0.10	0.10
Medium chain fatty acids ³	---	0.50
Total	100.00	100.00
Calculated analysis		
ME, kcal/lb	1,373	1,373
CP, %	21.14	21.14
SID Lys, %	1.12	1.12

¹ Treatments were arranged in 2×5 factorial design of diet form (meal vs. crumbled pellet) and 0.5% MCFA inclusion (control, C6, C8, C10, and C12).

² Composition per kg: 20 g iron, 40 g zinc, 40 g manganese, 4.5 g copper, 0.6 g iodine, and 0.06 g selenium. 3,080,000 IU vitamin A, 1,100,000 IU vitamin D₃, 6,600 IU vitamin E, 4.4 mg vitamin B₁₂, 330 mg menadione, 2,640 mg riboflavin, 2,640 mg d-pantothenic acid, and 11,000 mg niacin.

³ Medium chain fatty acids (Sigma Aldrich, St. Louis, MO) used in this experiment were guaranteed $\geq 98\%$ purity and included hexanoic (C6), octanoic (C8), decanoic (C10), and dodecanoic acid (C12).

Table 2. Effect of MCFA inclusion in mash or crumbled pellet diets on broiler performance through 18 days posthatch¹

	Mash					Crumbled pellet					SEM	Probability, <i>P</i> <		
	Control	C6	C8	C10	C12	Control	C6	C8	C10	C12		Form	MCFA	Interaction
BW, lb														
d 0	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.09	0.001	0.046	0.488	0.306
d 18	1.55	1.50	1.52	1.51	1.52	1.62	1.60	1.59	1.63	1.62	0.027	0.001	0.594	0.689
d 0 to 18														
BWG, lb	1.44	1.38	1.44	1.42	1.43	1.51	1.49	1.50	1.53	1.50	0.025	0.001	0.290	0.695
FI, lb	1.81	1.77	1.82	1.81	1.81	1.87	1.87	1.85	1.89	1.91	0.028	0.001	0.683	0.654
FCR	1.26	1.28	1.27	1.27	1.27	1.24	1.26	1.24	1.23	1.26	0.010	0.001	0.119	0.328

¹ At hatch, 400 male broilers (Cobb 500, Cobb-Vantress, Siloam Springs, AR) were placed in groups of 5 and reared to 18 days with 8 replicates per treatment. Dietary treatments were arranged in 2 × 5 factorial design of diet form (meal and crumbled pellet) and 0.5% MCFA inclusion (no inclusion, control; hexanoic acid, C6; octanoic acid, C8; decanoic acid, C10; and dodecanoic acid, C12).

^{a,b} Means within a row followed by different letter are significantly different ($P \leq 0.05$).

BWG = body weight gain. FI = feed intake. FCR = feed conversion ratio.