

Experiences with Metzenberg's method for preparing Neurospora DNA.

F. M. Nemeti

M. Schablik

G. Szabo

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Nemeti, F. M., M. Schablik, and G. Szabo (1982) "Experiences with Metzenberg's method for preparing Neurospora DNA.," *Fungal Genetics Reports*: Vol. 29, Article 10. <https://doi.org/10.4148/1941-4765.1639>

This Technical Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Experiences with Metzenberg's method for preparing Neurospora DNA.

Abstract

Experiences with Metzenberg's method for preparing Neurospora DNA.

Nemeti, F. M., M. Schablik and G. Szabo

Experiences with Metzenberg's method for
preparing Neurospora DNA.

Metzenberg and Baisch (1981 Neurospora Newsl. 28: 20) reported an easy method for preparing high molecular weight DNA from Neurospora crassa RL3-8 A. We confirmed their results: however, in our hands the nucleic acid prepared by Metzenberg's method contained large amounts of RNA and polysaccharides. To reduce contamination in our DNA preparation the original procedure was modified as follows: (a) digestion with RNase was extended to 3 hours; (b) final precipitation of DNA was performed with 0.54 volume of iso-propanol.

Since the product obtained in this way still contained equal amounts of DNA and RNA, as determined by the diphenylamine (Richards 1974 Anal. Biochem 57: 369) and phloroglucine (Dische and Barenfreund 1957 Biochim Biophys. Acta 23: 639) methods, respectively, an additional treatment with RNase was performed. The RNA content of the final product was less than 10%.

The DNA prepared either by the method of Metzenberg or according to the above modification contained about equal amounts of DNA and polysaccharides (w/w): the latter was determined with the anthron reagent (Roe 1955 J. Biol. Chem 212: 335).

The molecular weight of DNA was determined by agarose gel electrophoresis (Aaij and Borst 1972 Biochim Biophys. Acta 269: 192) with λ -phage DNA used as standard, and was found to be about 49 kilobase pairs (kb). About 5% of the product were fragments of lower molecular weight (between 23 and 49 kb).

In conclusion, although Metzenberg's method provides an easy way for obtaining high molecular weight DNA, the DNA prepared in that way is contaminated to a significant degree with RNA and polysaccharides.

Institute of Biology, University Medical School, H-4012 Debrecen, Hungary.