

Endogenous protease(s) in extracts of *Neurospora mycella* activate the exonuclease associated with a putative Rec-nuclease

M. J. Fraser
McGill University

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Fraser, M. J. (1976) "Endogenous protease(s) in extracts of *Neurospora mycella* activate the exonuclease associated with a putative Rec-nuclease," *Fungal Genetics Reports*: Vol. 23, Article 7. <https://doi.org/10.4148/1941-4765.1767>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Endogenous protease(s) in extracts of *Neurospora mycella* activate the exonuclease associated with a putative Rec-nuclease

Abstract

Activation of exonuclease by endogenous proteases

activate the **exonuclease** associated with a putative **Rec-nuclease**.

Fraser 1975 In P. C. Hanawalt and R.B. Setlow (Eds.) *Molecular mechanisms for repair of DNA*, part B, Plenum Publishing Corp., New York, p. 577). It was observed that the single-strand (s-) DNase activity of wild-type (74-OR23-IVA) had increased 3 to 4-fold and the double-strand (ds-) DNase had increased 20 to 25-fold over that in fresh extracts after storage at 0.4°C for more than two weeks. These activations occurred in 24 hr at room temperature (see Table I) and in 1-2 hr at 37°C. Both activities were subsequently reduced on further incubations.

Table I. Activation and inactivation of single-strand DNase (ssDNase) and double-strand DNase (dsDNase) activities in extracts of wild-type and uvs-3 by endogenous "serine proteinase"

Extract	Days at Room Temp.	Activity (units/ml)			
		ssDNase		dsDNase	
		-PMSF	+PMSF	-PMSF	+PMSF
Wild-type	0	87,88	98,83	3.0	2.2
"	1	336,340	106,101	66,64	1.9, 2.5
"	2	142	157	7.9	7.9
"	3	40	147	2.1	4.5
<u>uvs-3</u>	0	46.46	40,42	0.8	0.8
"	1	47	42,44	1.8, 1.5	1.1, 0.8
"	2	217	64	52.	2.1
"	3	105	66	15.	1.4

Aliquots of extracts of mycelia were put through 0.45 µm Millipore filters into sterile screw capped vials to avoid bacterial contamination. Crystals of solid PMSF were added to half of the vials (+PMSF). Room temperature averaged about 21°C.

37°C was inhibited by adding 2mg/ml serum albumin to the nuclease preparation. After pre-incubation without albumin, the nuclease activity remaining was found to be a single-strand specific endonuclease identical in properties with that described by Linn (see above).

A stable nuclease preparation has now been derived from the uvs-3 strain. When this was subjected to electrophoresis in 6M urea-polyacrylamide gels, a very acidic protein was recovered which had both ss-DNase and ds-DNase activities. When uvs-3 nuclease preparation was treated with 3-10 µg trypsin for 30 min at 37°C, the exonuclease was activated. It thus seems likely that the activation of the ds-DNase activity observed in extracts is due to the direct action of "serine proteinase" on the enzyme rather than due to the destruction of an inhibitor. It also seems possible that the uvs-3 strain is deficient in proteinase(s) which cause these cower-

■ ■ ■ Department of Biochemistry, McGill University, Montreal, Canada H3G 1Y6.

Recently, comparisons have been made of the single- and double-strand DNase activities of extracts of mycelia of wild-type, ultraviolet light-sensitive and putative DNase mutants of *Neurospora* (M. J.

The activation, and subsequent inactivation, of both DNase activities were much slower (but not prevented) in the presence of the "serine proteinase" inhibitor, phenylmethylsulfonyl fluoride (PMSF). In addition, both processes were reproducibly slower in extracts of the uvs-3 mutant (see Table I) which has been shown to have a phenotype similar to recA mutants of *E. coli* and may be altered in mitotic recombination (A. L. Schroeder 1970 *Molec. Gen. Genetics*, 107: 291-305). Fresh extracts of mycelia of uvs-3 have been found to have specific ss-DNase activities one-third that of wild-type. Most of the ss-DNase activity in mycelial extracts is associated with the single-strand specific endonuclease first described by Linn (1967 *Meth. in Enzymol.* 12A:247). When this activity was purified from log-phase wild-type mycelia, it was found to be associated with an exonuclease (Fraser and Tjeerde 1975 *Fed. Proc.* 34: 515) which is now known to have activity with ss-DNA and with linear, but not with circular, ds-DNA. The two activities comprise a putative Rec-nuclease. Purified wild-type nuclease preparations have been found to contain at least two proteinase activities, PMSF-sensitive and PMSF-insensitive (assayed using azoalbumin as substrate according to the method of Tomarelli et al. 1949 *J. Lab. Clin. Med.* 34: 428). Transient activations of the exonuclease activity have been observed in freshly purified wild-type nuclease preparations. Aging at 0-4°C or pre-incubating 1 hr at 37°C resulted in a preferential loss of exonuclease activity. The loss in activity at