

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 1312

1976

Feeding propionic acid-treated, flaked sorghum to finishing steers

Terry L. Gugle

G. Fink

Jack G. Riley

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Gugle, Terry L.; Fink, G.; and Riley, Jack G. (1976) "Feeding propionic acid-treated, flaked sorghum to finishing steers," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2715>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1976 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



K**S****U**

Feeding Propionic Acid-treated, Flaked Sorghum to Finishing Steers

Jack G. Riley, Terry Gogle, and Galen Fink

Summary

Fifty-four yearling Hereford steers averaging 805 pounds were used in an 86-day finishing trial to determine the effect of adding low levels of propionic acid to steam flaked sorghum grain. Three treatments were examined. Treatment one consisted of sorghum grain steam flaked twice weekly for a maximum of 4 days between flaking. Treatment two was the same as treatment 1 except that propionic acid was added into the discharge auger at 0.15% of the weight of the flakes. Treatment three was flaking once a week with propionic acid added at 0.25% of the weight of the flakes. Both acid levels significantly ($P < .05$) improved gain and efficiency.

Introduction

We, and commercial operators have observed that steam flaked grain appeared to lose quality and palatability when not fed the day it was processed. Even though flaked grain was dried to approximately 84% dry matter, it deteriorated noticeably when stored more than three days in warm humid weather. That prompted us to try propionic acid (less than 0.3% of flake weight) to maintain quality and to extend storage life of flaked grain.

Procedure

Fifty-four yearling Hereford steers averaging 805 lbs. were divided into 3 treatment groups for an 86-day trial. The three treatments were: (1) sorghum grain flaked twice per week for a maximum of 4 days storage between flaking; (2) sorghum grain flaked at the same intervals as treatment 1 with propionic acid added into the discharge auger at 0.15% of the weight of the flakes; (3) sorghum grain flaked once a week with propionic acid added at 0.25%. Composition of the rations on a dry matter basis was 80% steam flaked milo, 5% protein supplement, and 15% corn silage. The trial began November 12, 1974, and ended February 6, 1975. Individual weights were taken initially, at 28-day intervals, and at completion. Carcass data for each steer were collected at Wilson and Company, Kansas City, Mo.

Results

Results of trial 1 are shown in table 24.1. Treating with propionic acid significantly improved daily gain and efficiency of gain. Flaking twice a week and treating with 0.15% propionic acid reduced daily dry matter consumed so performance for this group was less than for the group getting milo flaked only once a week but treated with 0.25% propionic acid. This preliminary study indicates that adding propionic acid to steam flaked sorghum grain helps preserve feed quality so rate and efficiency of gain are improved.

Table 24.1 Effect of adding propionic acid to flaked sorghum grain on performance of finishing steers, Nov. 12, 1974 - Feb. 6, 1975 (86 days).

Item	Propionic acid		
	0	0.15%*	0.25%*
No. steers	18	18	18
Initial wt., lbs.	808	801.7	804.3
Final wt., lbs.	986.3	1009.8	1034.3
Gain, lbs.	178.3	208.1	230.0
A.D.G.	2.07 ^a	2.42 ^b	2.67 ^c
Dry matter/day, lbs.	19.00 ^b	17.44 ^a	19.48 ^b
Dry matter/lb gain, lbs.	9.18 ^a	7.21 ^b	7.30 ^b

*Of weight of flaked milo.

a,b,c Values in same line with different superscripts differ sign ($P < .05$).