

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 10 *Swine Day (1968-2014)*

Article 665

1996

Effects of fat and sodium bicarbonate on growth performance and stomach morphology in finishing pigs

S P. Sorrell

I H. Kim

Robert H. Hines

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

 Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Sorrell, S P.; Kim, I H.; Hines, Robert H.; Kennedy, G A.; Burnham, L L.; and Hancock, Joe D. (1996) "Effects of fat and sodium bicarbonate on growth performance and stomach morphology in finishing pigs," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 10. <https://doi.org/10.4148/2378-5977.6505>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1996 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Effects of fat and sodium bicarbonate on growth performance and stomach morphology in finishing pigs

Authors

S P. Sorrell, I H. Kim, Robert H. Hines, G A. Kennedy, L L. Burnham, and Joe D. Hancock

K

S

U

EFFECTS OF FAT AND SODIUM BICARBONATE ON GROWTH PERFORMANCE AND STOMACH MORPHOLOGY IN FINISHING PIGS

*S. P. Sorrell, J. D. Hancock, I. H. Kim,
R. H. Hines, G. A. Kennedy, and L. L. Burnham*

Summary

Pigs fed diets with soybean oil consumed less feed; grew more efficiently; and had greater last-rib backfat thickness, keratosis, and ulceration in their stomachs than pigs fed diets without soybean oil. Pigs fed diets with NaHCO_3 tended to eat more feed and had numerically greater ADG, but feed/gain and carcass measurements were not affected. NaHCO_3 decreased ulceration scores only for pigs fed diets without added fat.

(Key Words: Finishing Pigs, Fat, Sodium Bicarbonate, Ulcers.)

Introduction

Swine production in the United States has become very intensive. With this intensive production have come major changes in genetics; sometimes crowded pens; advanced feed processing procedures (e.g., fine grinding, pelleting, extruding, and expanding); and diet modifications (e.g., high fat) to maximize efficiency of gain. All of these factors have been implicated as contributors to the development of stomach lesions in finishing pigs. The resulting loss in growth performance and even death of afflicted pigs has prompted widespread investigation of methods to decrease the incidence and severity of stomach lesions.

Especially in the southeastern United States, anecdotal reports link crowded pens and high-fat diets with increased incidence of stomach lesions in pigs. In previous research here at KSU (1993 Swine Day Report), we observed that inclusion of 1% NaHCO_3 in simple corn-soy-based diets tended to reduce severity of gastric lesions in finishing pigs.

Thus, the objective of the experiment reported herein was to determine the effects of fat on growth performance and stomach lesions in finishing pigs and to evaluate NaHCO_3 as a modifying influence in development of stomach lesions for pigs fed diets with added fat.

Procedures

Two hundred and forty crossbred pigs were allotted to 16 pens with 15 pigs per pen (6 ft²/pig). Each pens was equipped with a self-feeder and nipple waterer to allowed ad libitum consumption of feed and water. The experiment was conducted as a 2 × 2 factorial with main effects of fat (none vs 6% added) and NaHCO_3 (none vs 1% added).

All pigs (Hampshire × Yorkshire × Duroc × Chester White crossbreds) were fed a corn-soybean meal-based diet formulated to .65% calcium and .55% phosphorus (Table 1). Lysine concentrations were .65% for the diets without added fat and .70% for the diets with added fat. The pigs were allotted to treatments on the basis of initial weight (avg of 100 lb), sex (eight barrows and seven gilts in each pen), and ancestry. The pigs were housed in a modified open-front building with half slatted and half solid, concrete floors.

The pigs were slaughtered when the average weight in the heaviest pen of a weight block reached 250 lb. At slaughter, carcass data were obtained, and the esophageal regions of the stomachs were scored for keratosis and ulceration. Response criteria were ADG, ADFI, G/F, backfat thickness, dressing percentage, FFLI, and scores for keratosis and ulceration in the stomach.

All data were analyzed as a randomized complete block design (initial weight as the blocking term) using the GLM procedure of SAS and with a 2×2 factorial arrangement of treatments. Pen was the experimental unit for all analyses.

Results and Discussions

No interactions ($P > .15$) occurred among the main effects of fat and NaHCO_3 for any of the growth or carcass data (Table 2). However, pigs fed diets with soybean oil consumed less feed and were more efficient than pigs fed diets without soybean oil ($P < .001$). These data are in agreement with numerous other reports that adding fat increases energy density of diets, reduces voluntary feed intake, and improves efficiency of growth. Pigs fed the fat-added diet had 1% greater dressing percentage ($P < .002$), but also had .08 in. more last rib fat thickness ($P < .006$).

The pigs ate more of the diets with 1% NaHCO_3 ($P < .06$), but rate and efficiency of gain were not affected ($P > .12$). Adding

NaHCO_3 to the diets did not affect carcass characteristics ($P > .15$).

Pigs fed soybean oil had greater keratinization ($P < .07$) and ulceration ($P < .05$) scores than pigs fed no soybean oil. Adding NaHCO_3 decreased ulceration scores, but only for pigs fed diets without added fat (fat \times NaHCO_3 interaction, $P < .01$). This response was actually the opposite of what was expected, i.e., that fat additions might aggravate stomach tissues and that NaHCO_3 could be of greatest benefit in those situations. To the contrary, lesion scores were actually slightly greater for pigs fed NaHCO_3 in the fat-added diets.

In conclusion, our data suggested that soybean oil enhanced growth performance but increased stomach keratosis and ulceration. Feeding NaHCO_3 did not affect growth performance (with only a trend for greater feed consumption) or carcass characteristics and prevented adverse changes in stomach morphology only for pigs fed diets without soybean oil.

Table 1. Diet Composition^a

Ingredient, %	No Soy Oil and No NaHCO_3	1% NaHCO_3	6% Soy Oil	1% NaHCO_3 & 6% Soy Oil
Corn	82.8	81.6	77.0	75.9
Soybean meal	14.6	14.8	14.2	14.3
Soy oil	-	-	6.0	6.0
NaHCO_3	-	1.0	-	1.0
Vit, Min, Antibio ^b	2.6	2.6	2.7	2.7
Lysine HCl	-	-	.1	.1
<u>Calculated analysis</u>				
Lys, %	.65	.65	.70	.70
ME, kcal/lb	1,508	1,492	1,611	1,595

^aFormulated to .65% Ca and .55% P.

^bSupplied 40 g/ton tylosin.

Table 2. Effects of Fat and Sodium Bicarbonate on Growth Performance, Carcass Characteristics, and Stomach Morphology in Finishing Pigs^a

Item	No Soy Oil & No NaHCO ₃	1% NaHCO ₃	6% Soy Oil	1% NaHCO ₃ & 6% Soy Oil	CV	Contrasts		
						1	2	3
ADG, lb	1.85	1.93	1.87	1.91	3.7	-. ^c	.12	-
ADFI, lb	5.86	6.10	5.54	5.64	2.7	.001	.06	-
F/G	3.18	3.17	2.96	2.96	3.0	.001	-	-
Backfat, in	1.14	1.15	1.24	1.20	3.7	.006	-	-
HCW, lb	186	186	189	189	.74	.002	-	-
Dressing, %	74.8	74.6	75.9	76.0	.8	.002	-	-
FFLI, %	47.1	46.9	46.2	46.6	.8	.01	-	-
Keratosisc	1.39	1.45	1.62	1.54	43	.07	-	-
Ulceration ^d	.72	.33	.66	.88	139	.05	-	.01

^aA total of 240 pigs (avg initial wt of 100 lb) were used.

^bContrasts were: 1) fat vs no fat; 2) NaHCO₃ vs no NaHCO₃; and 3) fat × NaHCO₃ interaction.

^cThe scoring system was: 0 = normal; 1 = mild; 2 = moderate; and 3 = severe keratosis.

^dThe scoring system was: 0 = normal; 1 = mild; 2 = moderate; and 3 = severe ulceration.

^eDashes indicate $P > .15$.



Swine Nutrition Graduate Students: Back row (L to R) Jim Smith, In-Ho Kim, Brandon Senne, Ioannis Mavromichalis, Robert Musser, Joe Loughmiller, LeAnn Johnston, Perry Sorrell; Front row (L to R) Carla Johnson, Jon Berstrom, Kittiporn Hongtrakul, Patrick O'Quinn, and Cao Hong. Not pictured, Chad Kerr and Ben Nessmith.