

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 10 *Swine Day (1968-2014)*

Article 1070

2014

Effects of an algae-modified montmorillonite clay on growth performance of nursery pigs fed diets contaminated with low levels of deoxynivalenol

Suzy Q. Fowler

Hyatt L. Frobose

Michael D. Tokach

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Fowler, Suzy Q.; Frobose, Hyatt L.; Tokach, Michael D.; DeRouchey, Joel M.; Dritz, Steven S.; Goodband, Robert D.; and Nelssen, Jim L. (2014) "Effects of an algae-modified montmorillonite clay on growth performance of nursery pigs fed diets contaminated with low levels of deoxynivalenol," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 10. <https://doi.org/10.4148/2378-5977.6910>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2014 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Effects of an algae-modified montmorillonite clay on growth performance of nursery pigs fed diets contaminated with low levels of deoxynivalenol

Authors

Suzy Q. Fowler, Hyatt L. Frobose, Michael D. Tokach, Joel M. DeRouchey, Steven S. Dritz, Robert D. Goodband, and Jim L. Nelssen

Effects of an Algae-Modified Montmorillonite Clay on Growth Performance of Nursery Pigs Fed Diets Contaminated with Low Levels of Deoxynivalenol¹

S.Q. Fowler, H.L. Frobose, M.D. Tokach, J.M. DeRouchey, S.S. Dritz², R.D. Goodband, and J.L. Nelssen

Summary

A total of 360 barrows (PIC 1050; initially 25.1 lb and 45 d of age) were used in a 21-d growth trial to evaluate the effects of an algae-modified montmorillonite clay (AMMC) on nursery pig performance when fed diets contaminated with low levels of deoxynivalenol (DON). Pigs were allotted to pens by weight, and pens were randomly assigned to 1 of 9 dietary treatments arranged in a 3 × 3 factorial with main effects of DON (0, 1.5 ppm, or 3.0 ppm) and AMMC inclusion (0, 0.17%, or 0.50%). There were 5 pigs per pen and 8 pens per treatment. Mycotoxin analyses were conducted on the main ingredients at NDSU³ and LDA Labs⁴, and the results were used in diet formulation. Naturally contaminated wheat (6.03 ppm DON) was used to produce diets with desired DON levels. No significant DON × AMMC interactions were observed during the entire study. Overall (d 0 to 21), increasing DON concentration in the diet decreased (1.22 vs. 1.10 vs. 1.07 lb; linear, $P < 0.001$) ADG and d-21 BW as a result of decreased ADFI (2.13 vs. 2.05 vs. 2.11 lb; quadratic, $P < 0.01$) and poorer feed efficiency (1.49 vs. 1.50 vs. 1.64; linear, $P < 0.001$). As expected, DON-related growth reductions were most marked from d 0 to 7 (15 to 22% lower) and least distinct in the final period, d 14 to 21 (5 to 6% lower). Incorporating AMMC at increasing levels had no effect on ADG, ADFI, feed efficiency, or final BW. Overall, the results of this study reinforce prior research showing that even low levels of DON significantly reduce nursery pig growth, but the addition of AMMC does not offset the deleterious effects of DON.

Keywords: deoxynivalenol, montmorillonite clay, nursery pigs, vomitoxin

Introduction

Deoxynivalenol (DON), also known as vomitoxin, is a mycotoxin commonly found in wheat, corn, and other cereal grains. During wet, cool periods of the growing season, *Fusarium graminearum* forms a head blight, producing DON as a secondary metabolite. Because DON occurs frequently and at toxicologically relevant concentrations and pigs are the most susceptible livestock species, it is one of the most important mycotoxins for swine producers to be aware of. At low concentrations (<1 ppm), DON causes decreased feed intake and upregulation of the immune system; at high levels (>10 ppm), it can cause complete feed refusal and vomiting. Deoxynivalenol remains challenging

¹ Appreciation is expressed to Olmix N.A. (Black River Falls, WI) for partial financial support.

² Department of Diagnostic Medicine/Pathobiology, College of Veterinary Medicine, Kansas State University.

³ North Dakota State University Veterinary Diagnostic Laboratory, Fargo, ND.

⁴ LDA Labs, Ploufragan, France.

for livestock producers because commonly used detoxifying agents such as bentonite clay and activated aluminosilicates, which are known to be effective against aflatoxins, are largely ineffective against DON. These additives may be ineffective against DON because the DON molecule is too large to become trapped within the clay matrix. However, it is hypothesized that an algae-modified montmorillonite clay (AMMC), which uses a patented process (Amadeite; Olmix, Brehan, France) to incorporate algae polysaccharides and expand the layers of the clay, may alleviate the growth reductions associated with feeding DON-contaminated grains. Therefore, the objective of this study was to evaluate the effects of currently recommended levels of AMMC on the growth performance of nursery pigs fed diets containing low levels of DON.

Procedures

The Kansas State University Institutional Animal Care and Use Committee approved the protocol used in this experiment. The trial was conducted at the K-State Segregated Early Weaning Facility in Manhattan, KS. A total of 360 barrows (PIC 1050; 25.1 ± 0.5 lb and 45 d of age) were used in a 21-d experiment with 8 replicate pens per treatment and 5 pigs per pen.

Pigs were shipped to the facility immediately postweaning and placed in 2 identical nurseries, each containing 40 pens (5 ft \times 5 ft). Upon arrival, pigs were placed in pens by weight and fed a common diet until approximately 25 lb.

On d 0, pigs were re-weighed and pens were randomly assigned to 1 of 9 dietary treatments in a 3×3 factorial arrangement with DON and AMMC inclusion as main effects (Table 1). The 9 dietary treatments consisted of 3 positive control (PC) diets without DON with 0, 0.17%, or 0.50% AMMC added, 3 low negative control (Low NC; 1.5 ppm DON) diets with 0, 0.17%, or 0.50% AMMC, and 3 high negative control (High NC; 3 ppm DON) diets with 0, 0.17%, or 0.50% AMMC. The AMMC that was supplied by Olmix N.A. (Black River Falls, WI) is typically recommended to be included at 0.17% and was added at the expense of corn in diet formulation. Diets exceeded NRC (2012⁵) nutrient requirements, and apart from the inclusion of DON and AMMC were formulated to be identical in nutrient composition.

Diets were manufactured in meal form at the K-State O. H. Kruse Feed Mill in Manhattan, KS. Hard red winter wheat naturally contaminated with DON was sourced (Table 2; 6.03 ppm DON) and incorporated into diets to achieve desired dietary DON levels. Prior to diet manufacturing, a total of 60 subsamples were collected from both a high-DON and DON-free wheat source. These samples were homogenized and split into duplicate samples, which were then sent for mycotoxin analysis at NDSU (North Dakota State University Veterinary Diagnostic Laboratory, Fargo, ND) and LDA Labs (Ploufragan, France). The lab at NDSU conducted an 18-component toxin screen using a combination of mass spectrometry, ELISA, and HPLC. LDA Labs performed a 43-component toxin screen using high-pressure liquid chromatography/mass spectrometry analyses. Due to concerns that high-DON wheat may also have a different amino acid profile than DON-free wheat, both wheat sources were analyzed for amino acid content (Table 3) at the University of Missouri Agricultural Experiment Station Chemical Laboratories (Columbia, MO), and diet formula-

⁵ NRC. 2012. Nutrient Requirements of Swine. 11th ed. Natl. Acad. Press, Washington, DC.

tion was adjusted to account for the differences. Following final diet manufacturing, diet samples were sent to NDSU and the University of Missouri for mycotoxin (Table 4) and proximate analysis (Table 5).

Average daily gain, ADFI, and F/G were determined by weighing pigs and determining feed disappearance on d 0, 7, 14, and 21 (Tables 6 and 7). Each pen contained a 4-hole, dry self-feeder and a nipple waterer to provide ad libitum access to feed and water.

Data were analyzed using the MIXED procedure of SAS (SAS Institute, Inc., Cary, NC) with the main effects and their interactions serving as fixed effects and barn as a random effect. Means were evaluated using preplanned linear and quadratic orthogonal contrasts. The coefficients for the unequally spaced linear and quadratic contrasts were derived using the IML procedure in SAS. Least squares means were calculated for each independent variable. Results were considered significant if P -values were ≤ 0.05 and tendencies if $0.05 > P \geq 0.10$.

Results and Discussion

Proximate and amino acid analysis of DON-contaminated wheat revealed protein and amino acid levels marginally but consistently higher than the DON-free wheat source. *Fusarium* pathogens are known to not only produce secondary metabolites such as mycotoxins but also alter the nutrient content of the affected grain source.⁶ Because these alterations in nutrient content are inconsistent, it is critical to account for these differences in diet formulation to assess the true impact of mycotoxin contamination on animal performance.

In the present study, analyzed DON concentrations in the naturally DON-contaminated wheat differed between NDSU (8.4 ppm) and LDA Labs (6.03 ppm). Although very low levels of several other mycotoxins were detected in the DON-contaminated wheat source, the levels observed were below concentrations necessary to elicit reductions in animal performance due to individual toxin exposure; however, the impact of interactive effects between toxins cannot be completely excluded. To ensure that final diet DON levels were adequate to achieve a DON-associated reduction in performance, the analysis from LDA Labs was used as the basis for diet formulation. Analyzed DON in the final diets revealed levels that were within 20% of the targeted DON level, averaging 1.73 and 3.20 ppm for the 1.5 and 3.0 ppm targets, respectively. Furthermore, proximate analyses of the final diets were generally in line with formulated levels.

Regarding growth performance, a tendency for a DON \times AMMC interaction (linear, $P < 0.10$) was observed for d 0 to 7, where increasing AMMC tended to improve ADG in PC and low-NC diets but tended to decrease ADG in high-NC diets. This interaction for ADG appeared to be driven by a tendency for a F/G interaction (linear, $P < 0.10$) in which increasing AMMC inclusion worsened feed efficiency in high-NC diets whereas feed efficiency remained similar in pigs fed PC and low-NC diets regardless of AMMC inclusion. Furthermore, a tendency for a DON \times AMMC interaction for feed efficiency (quadratic, $P < 0.10$) was observed from d 14 to 21 and overall (d 0 to 21),

⁶ Matthaues, K., S. Danicke, W. Vahjen, O. Simon, J. Wang, H. Valenta, K. Meyer, A. Strumpf, H. Ziesenib, and G. Flachowsky. 2004. Progression of mycotoxin and nutrient concentrations in wheat after inoculation with *Fusarium culmorum*. Arch. Anim. Nutr. 58:19–35.

where increasing AMMC in PC and low-NC diets worsened feed efficiency, whereas in high-NC diets increasing AMMC initially improved but subsequently worsened F/G at the 0.50% inclusion rate.

For the main effects of DON on nursery pig growth, results matched expected reductions in ADG, where pigs experienced the most deleterious effects of DON during the initial exposure period and seemed to attenuate somewhat over time. During d 0 to 7, ADG decreased (linear, $P < 0.001$) as DON was increased in diets, driven by both a decrease in feed intake (linear, $P < 0.001$) and a worsening of feed efficiency (linear, $P < 0.001$). These responses were expected because they are known to be associated with upregulation of the immune system during initial exposure. From d 7 to 14, reduced feed intake (quadratic, $P < 0.001$) continued to drive ADG lower (linear, $P < 0.001$) in pigs fed increasing concentrations of DON. However, increasing DON in the diets improved (quadratic, $P < 0.01$) feed efficiency, which may be attributed to improved nutrient utilization at such low feed intake. Nevertheless, from d 14 to 21, the previously observed negative effects of DON on nursery pig growth were less marked. Although there was a tendency for reduced ADG (linear, $P < 0.10$) as DON increased, ADFI and feed efficiency were similar across treatments. Overall (d 0 to 21), increasing DON concentration in nursery pig diets caused a reduction in ADG and BW (linear, $P < 0.001$), which was driven by lower ADFI (quadratic, $P < 0.01$) and poorer feed efficiency (quadratic, $P = 0.01$).

Although increasing DON in nursery pig diets resulted in the expected reductions in growth necessary to evaluate the potential effectiveness of a DON-detoxifying agent, the addition of AMMC at 0.17 and 0.50% did not affect ADG, ADFI, feed efficiency, or final pig BW during the experimental period.

In conclusion, although AMMC does not appear to offset the deleterious effects of low levels of DON in nursery pig diets, the results of this study indicate that minimizing DON contamination in swine diets is critical, because even low dietary levels (1.5 to 3.0 ppm) can result in growth performance reductions of 10% or greater. Correcting suboptimal field conditions to minimize DON concentration is impractical, so additional research is necessary to identify practical and effective methods to reduce the negative effects of deoxynivalenol postharvest.

Table 1. Formulated diet composition (as-fed basis)

Item	AMMC ² level:	Positive control (0 ppm DON ¹)			Low negative control (1.5 ppm DON)			High negative control (3.0 ppm DON)		
		None	0.17%	0.50%	None	0.17%	0.50%	None	0.17%	0.50%
Ingredient, %										
Corn		15.07	14.89	14.53	15.35	15.17	14.81	15.63	15.45	15.09
Soybean meal (46.5% CP)		31.58	31.60	31.62	31.25	31.26	31.29	30.92	30.93	30.96
Hard red winter (HRW) wheat		50.00	50.00	50.00	25.00	25.00	25.00	-	-	-
High-DON ³ HRW wheat		-	-	-	25.00	25.00	25.00	50.00	50.00	50.00
Monocalcium P, 21% P		1.05	1.05	1.05	1.05	1.05	1.05	1.05	1.05	1.05
Limestone		1.00	1.00	1.00	1.05	1.05	1.05	1.10	1.10	1.10
Salt		0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35
L-lysine HCl		0.33	0.33	0.33	0.33	0.33	0.33	0.33	0.33	0.33
DL-methionine		0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
L-threonine		0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13
Vitamin premix with phytase ⁴		0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Trace mineral premix		0.15	0.15	0.15	0.15	0.15	0.15	0.15	0.15	0.15
AMMC		-	0.17	0.50	-	0.17	0.50	-	0.17	0.50
Total		100	100	100	100	100	100	100	100	100
Calculated composition, %										
Standardized ileal digestible (SID) amino acids, %										
Lysine		1.28	1.28	1.28	1.28	1.28	1.28	1.28	1.28	1.28
Isoleucine:lysine		64	64	64	65	65	65	65	65	65
Leucine:lysine		118	118	117	118	118	118	117	117	117
Methionine:lysine		31	31	31	31	31	31	31	31	31
Met & Cys:lysine		57	57	57	57	57	57	57	57	57
Threonine:lysine		63	63	63	63	63	63	63	63	63
Tryptophan:lysine		21.2	21.2	21.2	20.7	20.7	20.7	20.3	20.3	20.2
Valine:lysine		68	68	68	69	69	69	70	70	70
Total lysine, %		1.43	1.43	1.43	1.43	1.43	1.43	1.43	1.43	1.43
ME, kcal/lb		1,444	1,442	1,436	1,443	1,441	1,436	1,443	1,440	1,435
SID lysine:ME, g/Mcal		4.02	4.03	4.04	4.02	4.03	4.04	4.02	4.03	4.05
CP, %		22.7	22.7	22.6	22.6	22.6	22.6	22.6	22.6	22.6
Ca, %		0.68	0.68	0.68	0.69	0.69	0.69	0.71	0.71	0.71
P, %		0.68	0.68	0.68	0.69	0.69	0.69	0.71	0.71	0.70
Available P, %		0.50	0.50	0.50	0.51	0.51	0.51	0.51	0.51	0.51
DON, ppm		<0.5	<0.5	<0.5	1.5	1.5	1.5	3.0	3.0	3.0

¹ Deoxynivalenol (DON).

² AMMC algae-modified montmorillonite clay product (Olmix, Brehan, France).

³ Analyzed DON concentration in wheat was 6.03 ppm at LDA Laboratories (Ploufragan, France).

⁴ Phyzyme 600 (Danisco Animal Nutrition, St. Louis, MO) provided 750 phytase units (FTU) phytase/kg and 0.13% available P released.

Table 2. Mycotoxin analysis of basal ingredients

Item, ppm	Ground corn	Hard red winter wheat	
		DON-free	High-DON ¹
NDSU ²			
DON	<0.50	<0.50	8.40
LDA Labs ³			
DON	--- ⁴	---	6.03
De-epoxy DON	---	---	0.02
15-O-Acetyl DON	---	---	0.07
3-Acetyl DON	---	---	0.03
Zearalenone	---	---	0.02
HT-2 Toxin	---	---	0.02
Ergocryptin	---	---	0.08
Ergosin	---	---	0.02
Tenuazonic acid	---	---	0.05

¹Deoxynivalenol (DON).

²North Dakota State University Veterinary Diagnostic Laboratory (Fargo, ND). Samples were sent for 18-component mycotoxin analysis and analyzed using a variety of mass spectrometry, ELISA, and HPLC methods. Included in the table are mycotoxins found at levels above detection limits.

³LDA Labs (Ploufragan, France). Samples analyzed using a 43-component toxin screen using liquid chromatography/mass spectrometry analysis methods. Included in the table are mycotoxins found at levels above detection limits.

⁴(---) indicates samples were not tested.

Table 3. Nutrient analysis of basal ingredients (as-fed basis)¹

Item, %	Hard red winter wheat	
	DON-free	High-DON ²
Moisture	9.14	10.20
Crude protein	11.80	12.20
ADF	3.20	2.50
NDF	8.10	10.30
Crude fat	1.80	2.00
Ash	1.81	1.84
Ca	0.07	0.06
P	0.38	0.44
Amino acid analysis		
Lysine	0.40	0.44
Isoleucine	0.41	0.47
Leucine	0.84	0.87
Methionine	0.21	0.22
Cysteine	0.27	0.27
Threonine	0.36	0.38
Tryptophan	0.15	0.17
Valine	0.57	0.47

¹ Samples were analyzed at the University of Missouri Agricultural Experiment Station Chemical Laboratories (Columbia, MO).

² Deoxynivalenol (DON).

Table 4. Mycotoxin analysis of experimental diets (as-fed basis)¹

Item	AMMC ³ level:	Positive control (0 ppm DON ²)			Low negative control (1.5 ppm DON)			High negative control (3.0 ppm)		
		None	0.17%	0.50%	None	0.17%	0.50%	None	0.17%	0.50%
DON, ppm		<0.5	<0.5	<0.5	1.7	1.8	1.7	3.4	2.7	3.5

¹ Diet samples were analyzed at North Dakota State University Veterinary Diagnostic Laboratory (Fargo, ND). An 18-component mycotoxin analysis was conducted using a variety of mass spectrometry, ELISA, and HPLC methods. Included in the table are mycotoxins found at levels above detection limits.

² Deoxynivalenol (DON).

³ AMMC. Algae-modified montmorillonite clay (Olmix, Brehan, France).

Table 5. Nutrient analysis of experimental diets (as-fed basis)¹

Item, %	AMMC level ³ :	Positive control (0 ppm DON ²)			Low negative control (1.5 ppm DON)			High negative control (3.0 ppm)		
		None	0.17%	0.50%	None	0.17%	0.50%	None	0.17%	0.50%
Moisture		9.77	9.60	9.93	10.04	9.77	9.89	9.97	9.66	9.75
Crude protein		24.9	23.8	24.2	23.4	23.2	23.3	23.5	23.7	23.5
ADF		2.6	2.4	2.1	2.7	3.5	2.2	2.5	2.6	2.6
NDF		7.6	7.0	7.5	7.6	8.0	7.4	6.8	7.2	7.2
Crude fat		2.4	2.6	2.5	2.6	2.9	2.6	2.6	2.8	2.7
Ash		5.14	5.31	5.53	5.53	5.61	5.57	5.65	5.73	5.96
Ca		0.71	0.81	0.82	0.86	0.83	0.76	0.87	0.83	0.89
P		0.74	0.67	0.68	0.69	0.69	0.71	0.69	0.71	0.74

¹ Samples were analyzed at the University of Missouri Agricultural Experiment Station Chemical Laboratories (Columbia, MO).

² Deoxynivalenol (DON).

³ AMMC. Algae-modified montmorillonite clay (Olmix, Brehan, France).

Table 6. Effects of an algae-modified montmorillonite clay (AMMC) on growth performance of nursery pigs fed diets contaminated with low levels of deoxynivalenol (DON)¹

Probability, <i>P</i> <													
DON level ² :		Positive control (0 ppm DON)			Low negative control (1.5 ppm DON)			High negative control (3.0 ppm DON)			SEM	DON × AMMC	
Item	AMMC ³ level:	None	0.17%	0.50%	None	0.17%	0.50%	None	0.17%	0.50%		Linear	Quadratic
d 0 to 7													
	ADG, lb	0.92	0.86	0.94	0.77	0.75	0.81	0.75	0.74	0.64	0.041	0.053	0.849
	ADFI, lb	1.38	1.38	1.40	1.23	1.24	1.29	1.22	1.21	1.16	0.043	0.235	0.803
	F/G	1.49	1.61	1.50	1.62	1.67	1.62	1.66	1.66	1.84	0.060	0.065	0.766
d 7 to 14													
	ADG, lb	1.27	1.26	1.32	1.18	1.08	1.16	1.07	1.06	1.12	0.071	0.987	0.315
	ADFI, lb	1.56	1.68	1.63	1.52	1.42	1.48	1.27	1.35	1.30	0.118	0.258	0.843
	F/G	1.23	1.34	1.23	1.29	1.31	1.29	1.18	1.26	1.14	0.087	0.332	0.676
d 14 to 21													
	ADG, lb	1.48	1.52	1.44	1.47	1.38	1.33	1.41	1.42	1.41	0.046	0.527	0.277
	ADFI, lb	2.12	2.11	2.15	2.05	2.06	2.03	2.16	2.03	2.14	0.187	0.851	0.424
	F/G	1.43	1.39	1.50	1.40	1.51	1.54	1.53	1.43	1.52	0.132	0.391	0.063
d 0 to 21													
	ADG, lb	1.23	1.21	1.23	1.14	1.07	1.10	1.08	1.07	1.06	0.032	0.618	0.268
	ADFI, lb	1.78	1.81	1.86	1.65	1.63	1.67	1.79	1.68	1.75	0.057	0.300	0.884
	F/G	1.45	1.50	1.51	1.45	1.53	1.52	1.67	1.57	1.67	0.063	0.570	0.090
Pig weight, lb													
	d 0	25.1	25.2	25.2	25.1	25.1	25.2	25.2	25.1	25.1	0.53	0.965	0.996
	d 7	31.6	31.2	31.8	30.5	30.4	30.8	30.4	30.3	29.6	0.68	0.415	0.936
	d 14	40.5	40.0	41.0	38.8	37.9	39.0	37.9	37.7	37.4	0.86	0.523	0.662
	d 21	50.8	50.6	51.1	49.1	47.5	48.3	47.7	47.6	47.3	1.05	0.740	0.488

¹ A total of 360 barrows (PIC 1050; initially 25.1 lb and approximately 45 d of age) were used in a 21-d experiment with 8 replicate pens per treatment and 5 pigs per pen. All diets were fed in meal form.

² Denotes formulated levels. Wheat naturally contaminated with DON (6.03 ppm) was used to incorporate DON into diets at desired concentrations.

³ AMMC. Algae-modified montmorillonite clay (Olmix, Brehan, France).

Table 7. Main effects of deoxynivalenol (DON) and algae-modified montmorillonite clay (AMMC) on nursery pig performance¹

Item	Formulated DON level ² , ppm				AMMC ³ level, %				Probability, <i>P</i> <			
									DON		AMMC	
	0	1.5	3.0	SEM	None	0.17%	0.50%	SEM	Linear	Quadratic	Linear	Quadratic
d 0 to 7												
ADG, lb	0.91	0.77	0.71	0.026	0.81	0.78	0.80	0.026	<0.001	0.188	0.747	0.353
ADFI, lb	1.39	1.25	1.19	0.021	1.27	1.28	1.28	0.021	<0.001	0.181	0.789	0.979
F/G	1.53	1.63	1.72	0.057	1.59	1.65	1.65	0.057	<0.001	0.848	0.261	0.396
d 7 to 14												
ADG, lb	1.28	1.14	1.08	0.060	1.17	1.13	1.20	0.060	<0.001	0.218	0.747	0.148
ADFI, lb	1.62	1.48	1.30	0.091	1.45	1.48	1.47	0.091	0.764	<0.001	0.650	0.293
F/G	1.26	1.30	1.20	0.050	1.23	1.30	1.22	0.050	<0.001	<0.01	0.616	0.916
d 14 to 21												
ADG, lb	1.48	1.39	1.41	0.027	1.45	1.44	1.39	0.027	0.087	0.103	0.652	0.935
ADFI, lb	2.13	2.05	2.11	0.178	2.11	2.06	2.11	0.178	0.754	0.166	0.545	0.370
F/G	1.44	1.48	1.49	0.126	1.46	1.44	1.52	0.126	0.199	0.592	0.741	0.310
d 0 to 21												
ADG, lb	1.22	1.10	1.07	0.020	1.15	1.12	1.13	0.020	<0.001	0.053	0.644	0.292
ADFI, lb	1.82	1.65	1.74	0.038	1.74	1.71	1.76	0.038	0.357	<0.01	0.629	0.233
F/G	1.49	1.50	1.64	0.050	1.52	1.53	1.57	0.050	<0.001	0.010	0.120	0.790
Pig weight, lb												
d 0	25.1	25.1	25.1	2.55	25.1	25.1	25.2	2.55	0.999	0.979	0.968	0.998
d 7	31.5	30.6	30.1	3.23	30.8	30.6	30.7	3.23	0.013	0.583	0.922	0.709
d 14	40.5	38.5	37.7	4.35	39.0	38.5	39.1	4.35	<0.001	0.376	0.772	0.406
d 21	50.8	48.3	47.6	5.38	49.2	48.6	48.9	5.38	<0.001	0.220	0.789	0.510

¹ A total of 360 barrows (PIC 1050; initially 25.1 lb and approximately 45 d of age) were used in a 21-d experiment with 24 replicate pens per treatment and 5 pigs per pen. All diets were fed in meal form.

² Denotes formulated levels. Wheat naturally contaminated with DON (6.03 ppm) was used to incorporate DON into diets at desired concentrations.

³ AMMC. Algae-modified montmorillonite clay (Olmix, Brehan, France).