

# Kansas Agricultural Experiment Station Research Reports

---

Volume 0  
Issue 12 *Keeping up with Research*

Article 43

---

1974

## Feeding Holstein Bulls and Steers for Beef Production (1974)

Leslie J. Chyba

Fred W. Boren

Follow this and additional works at: <https://newprairiepress.org/kaesrr>

---

### Recommended Citation



Chyba, Leslie J. and Boren, Fred W. (1974) "Feeding Holstein Bulls and Steers for Beef Production (1974)," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 12. <https://doi.org/10.4148/2378-5977.7280>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1974 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



KEEPING UP WITH RESEARCH 12  
Contribution No. 42  
November 1974

FEEDING HOLSTEIN BULLS AND STEERS FOR BEEF PRODUCTION



AGRICULTURAL EXPERIMENT STATION  
Kansas State University, Manhattan  
Floyd W. Smith, Director  
  
SOUTHEAST KANSAS BRANCH  
Mound Valley  
Fred W. Boren, Superintendent

## Feeding Holstein Bulls and Steers for Beef Production

Leslie J. Chyba, Beef Scientist  
Fred W. Boren, Station Superintendent

Research has shown that bulls gain faster, are more efficient, and produce leaner carcasses than steers, thus creating considerable interest in fattening young bull calves. Advantages are capitalizing on natural growth-promoting hormones of intact animals, and eliminating stress from castration. Grading standards for "bullock beef" (under 24 months old) should insure markets for such animals.

This trial included 12 Holsteins, six yearling steers and six bull calves. They were slaughtered at approximately the same weight. The younger bull calves had to be fed longer because they weighed less at starting than the steers did. Both lots were fed twice daily a ration of 80% rolled milo, 15% brome hay, and 5% soybean meal based supplement. All were weighed every 28 days.

Results indicate that bulls gained faster and were more efficient feed converters than steers (Table 1). Carcass data clearly show bulls produced leaner carcasses having less kidney knob and back fat cover, larger loin eyes, and much lower marbling scores.

Table 1. Performance of young bulls and steers compared.

	Holsteins	
	Bulls	Steers
No. of animals	6	6
Avg. initial wt., lbs.	388.2	628.5
Avg. final wt., lbs.	963.8	991.5
Avg. wt. gain, lbs.	575.6	363.0
Avg. daily gain, lbs.	2.57	2.05
Days on feed	244	177
<u>Avg. daily ration, lbs.</u>		
Grain	16.15	18.25
Hay	2.58	3.28
Supplement	1.00	1.00
Total daily dry matter intake, lbs.	19.73	22.53
<u>Carcass data</u>		
Hot wt., lbs.	542.3	580.0
Dressing %	56.4	57.9
Marbling <sup>a</sup>	8.0	14.2
Conformation <sup>b</sup>	17.8	17.2
Kidney knob, %	2.0	2.5
Loin eye, sq. in.	11.27	8.8
Back fat thickness, in.	.06	.11
USDA grade <sup>c</sup>	17.2	18.3
Yield grade <sup>d</sup>	1.3	2.6

a. 8 = traces, 14.2 = small.

b. 16 = low good, 17 = good, 18 = high good.

c. 17 = middle good.

d. Yield grade 1 to 5, with 1 most desirable.

Information in this report is for farmers, producers, colleagues, industry cooperators, and other interested persons. It is not a recommendation or endorsement as it is not yet backed by enough research.

Contribution no. 42, Southeast Kansas Branch Experiment Station, Mound Valley, Kansas Agricultural Experiment Station, Kansas State University.

Publications and public meetings by the Kansas Agricultural Experiment Station are available and open to the public regardless of race, color, national origin, sex, or religion.