

New chromosome rearrangements from conidia of *Neurospora* wild type OR23-1VA

David D. Perkins
Stanford University

J. A. Kinsey
University of Kansas Medical School

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Perkins, D. D., and J.A. Kinsey (1993) "New chromosome rearrangements from conidia of *Neurospora* wild type OR23-1VA," *Fungal Genetics Reports*: Vol. 40, Article 27. <https://doi.org/10.4148/1941-4765.1418>

This Regular Paper is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in *Fungal Genetics Reports* by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

New chromosome rearrangements from conidia of *Neurospora* wild type OR23-1VA

Abstract

In experiments of Perkins, Kinsey, Asch and Frederick (1993 *Genetics* 134:729-736), new chromosome rearrangements were recovered with a frequency of 10% or more among mitotically stable transformants of *am* strains, whereas rearrangements were rare (< 1%) in cultures from untransformed regenerated protoplasts of the same strains.

New chromosome rearrangements from conidia of *Neurospora* wild type OR23-1VA.

D.D. Perkins and J.A. Kinsey - Department of Biological Sciences, Stanford University, Stanford CA 94305-5020 and Department of Microbiology, Molecular Genetics and Immunology, University of Kansas Medical School, Kansas City KS 66160-7420

In experiments of Perkins, Kinsey, Asch and Frederick (1993 *Genetics* 134:729-736), new chromosome rearrangements were recovered with a frequency of 10% or more among mitotically stable transformants of *am* strains, whereas rearrangements were rare (< 1%) in cultures from untransformed regenerated protoplasts of the same strains. Rearrangement frequencies have also been determined for Oak Ridge wild types, although these were not the strains used for transformation and were therefore not appropriate as controls. Isolates were examined from regenerated untransformed protoplasts or from unprotoplasted conidia, with the results shown in Table 1. Surprisingly, 14 translocations were found in 278 isolates (5%) in five experiments with strain OR23-1VA (FGSC 2489), which has been used for many years as a standard laboratory wild type. Several apparently spontaneous translocations have been recovered in progeny from OR23-1VA during this time, but there has been no reason to suspect instability of the magnitude observed here. The translocations isolated from OR23-1VA all have different breakpoints. Stocks of OR23-1VA from three different independently maintained sources were used in the various series.

In contrast, only one translocation was found in 520 isolates (0.2%) with the other untransformed strains that were tested. No rearrangement was detected in 57 regenerated protoplasts from ORS-6a (FGSC 2490), and none in 196 from strain QNS-6 *nit-2 leu-3 caf-1 at a* (FGSC 5381; described by Perkins et al. 1986 *Genetics* 114:791-817). Previously, when the same QNS-6 strain was crossed with several normal sequence strains, 714 single-ascospore cultures were tested without finding a single rearrangement (V. C. Pollard and Perkins, unpublished).

The Oak Ridge strain of *a* mating type, ORS-6a, is a product of 13 successive backcrosses to OR23-1VA or to its vegetative precursor OR23-1A (Newmeyer, Perkins, and Barry 1987 *Fungal Genet. Newsl.* 34:46-51). The OR *A* and *a* wild types should therefore be nearly isogenic (except for regions close to mating type), yet the *A* strain seems markedly more prone to rearrangement. The cause of instability in OR23-1VA is unknown.

TABLE 1

Chromosome rearrangements recovered from single conidial isolates in the absence of transformation

| Strain | Experiment | No. of isolates tested | No. of confirmed rearrangements | Rearrangements recovered |
|--|------------|------------------------|---------------------------------|---|
| <i>am₆af^sa</i> | UK17 | 122 | 1 | <i>T(I;VI)UK17-51</i> |
| <i>am₁₃₂int^l A</i> | UK18 | 143 | 0 | |
| QNS-6 a | QNS | 198 | 0 | |
| ORS-6a | UK16 | 57 | 0 | |
| OR23-1VA | UK7 | 27 | 2 | <i>T(IIIR;VR)UK7-7 mei</i> , <i>T(V;VI)UK7-11</i> |
| " | UK8 | 36 | 2 | <i>T(IIIR;IL)UK8-18</i> , <i>T(I;IV;VR)UK8-21</i> , |
| " | UK14 | 58 | 5 | <i>T(VIR→VL)UK14-1</i> , <i>T(I;VI)UK14-2</i> , <i>T(II;VR)UK14-3</i> , <i>T(III;IV;VII)UK14-5</i> , <i>T(I;II;VI)UK14-7</i> |
| " | UK15 | 57 | 2 | <i>T(I;V)UK15-1</i> , <i>T(VI;VII)UK15-2</i> |
| " | UK19 | 100 | 3 | <i>T(IV;VII)UK19-4</i> , <i>T(IL;VI)UK19-37</i> , <i>T(VI;VII)UK19-65</i> |

UK17 and UK18 are untransformed control experiments from Perkins, Kinsey, Asch, and Frederick (1993 Genetics 134:729-736). Tested cultures in all the experiments were derived from single untransformed, regenerated protoplasts except for experiment UK19, in which they were from untransformed, unprotoplasted single conidia. Cultures were purified by serial transfer. Breakpoints are different in the two I;VI translocations.