

Isozyme variation in natural populations

M. M. Reddy
McMaster University

S. F.H. Threlkeld
McMaster University

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Reddy, M. M., and S.F. Threlkeld (1971) "Isozyme variation in natural populations," *Fungal Genetics Reports*: Vol. 18, Article 7. <https://doi.org/10.4148/1941-4765.1886>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Isozyme variation in natural populations

Abstract

Isozyme variation in natural populations

Reddy, M. M. and S. F. H. Threlkeld. Isozyme variation in natural populations of *Neurospora*.

Eight heterothallic (P384, P385, P406, P407, P413, P419, P438, P439) and three homothallic strains (P388, P404, P435) of *Neurospora* were obtained from D. D. Perkins' Florida collection. Mycelial extracts from these strains were subjected to acrylamide and starch gel electrophoresis. Out of ten enzymes examined, electrophoretic variation was observed only for esterases. The sites of esterase activity were numbered from 1 to 4 in order of rate of movement towards the anode, with site 1 being the fastest.

Of the eight heterothallic strains, six (P384, P385, P406, P407, P413, P419) had esterase site 1 and two strains (P438, P439) had both esterase sites 1 and 2. Of the three homothallic strains, two (P388, P436) had esterase sites 3 and 4 and the third (P404) had esterase site 2. Amylase, aminopeptidase, o-glycerophosphate dehydrogenase, 6-phosphogluconate dehydrogenase and indophenol oxidase showed one site of activity. Acid phosphatase showed activity at two sites and lactate dehydrogenase, peroxidase and glucose-6-phosphate dehydrogenase showed activity at three sites for all the strains. The absence of electrophoretic variation for these enzymes suggests that selection may have been operating against enzyme variants resulting in stabilization of the enzyme genotype of isolated populations in nature.

We would like to thank D. D. Perkins for kindly providing the strains. — — — Department of Biology, McMaster University, Hamilton, Ontario, Canada.