

Linkage data on phen and tyr mutants

A. A. El-Eryani

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

El-Eryani, A. A. (1968) "Linkage data on phen and tyr mutants," *Fungal Genetics Reports*: Vol. 13, Article 8.
<https://doi.org/10.4148/1941-4765.1941>

This Linkage Data is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Linkage data on phen and tyr mutants

Abstract

Linkage data on *phen* and *tyr* mutants

El-Eryani, A. A. Linkage data on phen

and tyr mutants.

The phenylalanine requiring mutants phen-2 (E5212) and phen-3 (Y 16329) are bath alleles at the same locus and are in the right arm of linkage group III at 2.2 map units to the left of tyr-1 (Y6994).

The tyrosine-requiring strain tyr(NH)₁₀₀spans also to tryptophan and leucine just as phen-1 does.

— — — Department of Biology, Yale University, New Haven, Connecticut 06520.